

Teen Pregnancy Rates by County 2009 – 2013
Prepared by LaShanda Eller, MPH. Oregon Public Health Division

Pregnancy Rate Per 1,000 Females ages 15-19 by County of Residence

County of Residence	2009	2010	2011	2012	2013
State Total	45.27	39.25	35.95	32.24	29.21
Baker	46.05	35.95	37.77	37.64	42.18
Benton	17.12	13.12	12.65	12.56	11.63
Clackamas	32.76	27.65	22.89	21.99	20.34
Clatsop	41.33	38.25	31.51	32.48	41.91
Columbia	41.16	41.23	28.21	21.79	30.18
Coos	45.86	52.25	45.39	45.25	36.84
Crook	45.60	43.78	43.47	20.47	31.35
Curry	34.42	40.54	32.25	46.61	28.26
Deschutes	39.57	36.90	33.93	26.64	25.15
Douglas	51.61	48.74	48.21	48.04	42.12
Gilliam	**	**	**	**	**
Grant	65.93	74.86	32.93	**	**
Harney	47.82	43.10	44.64	**	41.66
Hood River	41.77	32.29	31.00	34.30	24.03
Jackson	50.89	41.56	41.63	37.56	34.75
Jefferson	76.10	77.66	58.82	48.92	62.11
Josephine	42.71	50.89	47.41	39.16	35.14
Klamath	56.49	58.63	57.15	54.95	49.18
Lake	25.31	**	**	46.29	**
Lane	32.97	29.60	27.39	24.35	23.57
Lincoln	63.39	54.61	51.55	40.63	28.82
Linn	56.41	42.69	38.47	40.39	32.65
Malheur	77.48	71.16	73.28	54.02	55.1
Marion	63.18	55.08	47.93	40.29	35.94
Morrow	61.88	52.00	57.87	28.70	27.43
Multnomah	55.00	43.74	41.10	35.78	32.18
Polk	37.50	28.50	26.49	25.09	18.33
Sherman	**	**	**	**	**
Tillamook	43.73	41.92	48.46	42.37	33.33
Umatilla	67.90	66.18	69.31	57.60	54.87
Union	43.24	38.37	39.41	24.73	26.28
Wallowa	**	**	**	**	**
Wasco	46.99	43.30	51.72	41.48	39.94
Washington	36.06	31.16	27.47	26.89	23.78
Wheeler	**	**	**	**	**
Yamhill	42.11	32.91	28.57	29.89	24.25

**Rates and percentages based on 5 or fewer events are not reported.

Department of Health & Human Services, Office of Adolescent Health

Evidence-Based TPP Programs Eligible for Replication*

Evidence-Based TPP Programs are Implementation Ready and Eligible for Replication with OAH Funding							
Program Model	Program Type	Implementation Setting	Implementation Length	Population	Outcomes	Duration of Outcomes	Evidence Review Rating
Aban Aya Youth Project	Sexuality education	In-school grades 5-8 Community-based	16-21, 45-min lessons each year in grades 5-8	African American youth in grades 5-8	Recent sexual activity	Post-intervention (Spring 8 th grade) for boys only	Moderate
Adult Identity Mentoring (Project AIM)	Youth development	In-school middle school After-school Community-based	12 sessions, 50-min each	Low-income youth ages 11-14	Recent sexual activity	3-months post intervention	High
All4You!	Programs for youth in alternative schools	In-school or non-school alternative education settings	14 sessions ranging from 70-140 minutes each; service learning component	Alternative high school students ages 14-18	Frequency of sexual activity Contraceptive use and consistency	Six months after baseline	High
Be Proud! Be Responsible!	Sexuality education	In-school high school After-school Community-based	6 sessions, 60-min each	Youth ages 13-18	Recent sexual activity Number of sexual partners Frequency of sexual activity Contraceptive use and consistency	Ranges, 3-12 months post intervention	High
Be Proud! Be Responsible! Be Protective!	Program for Pregnant and Parenting Girls	In-school high school After-school Community-based	8 sessions, 60-min each	Pregnant and parenting girls in grades 7-12	Number of sexual partners	12-months post intervention	High

Evidence-Based TPP Programs are Implementation Ready and Eligible for Replication with OAH Funding

Program Model	Program Type	Implementation Setting	Implementation Length	Population	Outcomes	Duration of Outcomes	Evidence Review Rating
Becoming a Responsible Teen (BART)	Sexuality education	Community-based	8 sessions, 1.5-2 hours each	Youth ages 14-18	Recent sexual activity	12-months post intervention	High
Children's Aid Society (CAS)-Carrera Program	Youth development	After school program Community-based In-school	Daily 2-3 hour sessions for 7 years	Disadvantaged youth ages 11-12	Recent sexual activity Pregnancy or birth	Three years after program start for girls only	High
Cuidate!	Sexuality education	In-school high school After-school Community-based	6, 1-hour lessons	Latino youth ages 13-18	Recent sexual activity Number of sexual partners	Ranges, 3-12 months post intervention	High
Draw the Line/Respect the Line	Sexuality education	In-school middle school After-school Community-based	5 lessons in 6 th grade, 7 lessons in 7 th grade, 7 lessons in 8 th grade	Youth in grades 6-8	Sexual initiation or abstinence Recent sexual activity	12-months post intervention for boys only	High
Families Talking Together (FTT)	Program for Families	Clinic-based Community-based	11 modules; # sessions varies depending on program delivery	Parents of Latino & African American adolescents ages 10-14	Sexual initiation or abstinence Frequency of sexual activity	9-months post intervention	High
FOCUS	Sexuality education	In-school high school & college Community	4, 2-hour sessions	Women ages 16 and older	Number of sexual partners	11-months post intervention	High
Health Improvement Project for Teens (HIP Teens)	Sexuality education	After-school Community-based	4, 120-minute sessions; 2, 90-minute booster sessions	Low income, urban, sexually active girls ages 15-19	Recent sexual activity Number of partners Frequency of sexual activity Contraceptive use	6-months post intervention	High

Evidence-Based TPP Programs are Implementation Ready and Eligible for Replication with OAH Funding

Program Model	Program Type	Implementation Setting	Implementation Length	Population	Outcomes	Duration of Outcomes	Evidence Review Rating
Heritage Keepers Abstinence Education	Abstinence education	In-school (7-9 th grade)	5, 90-min sessions or 10, 45-min sessions	Youth in grades 7-9	Sexual initiation or abstinence	12-months post intervention	Moderate
HORIZONS	Sexuality education	Clinic-based	2, 4-hour sessions; 4, 15-minute telephone follow-ups	Females ages 15-21	Contraceptive use and consistency Sexually transmitted infections or HIV	12-months post intervention	High
It's Your Game: Keep it Real	Sexuality education	In-school middle school (grades 7-8) After-school	12, 50-min lessons in 7 th grade and 12, 50-min lessons in 8 th grade	Youth in grades 7-8; allowable for youth in 6 th grade	Sexual initiation or abstinence Frequency of sexual activity Contraceptive use and consistency	12-months post intervention	Moderate
Making a Difference!	Abstinence education	In-school high school After-school Community-based	8, 1-hour sessions	Youth ages 11-13	Recent sexual activity	3-months post intervention	High
Making Proud Choices!	Sexuality education	In-school high school After-school Community-based	8, 1-hour sessions	Youth ages 11-13; older teens allowable; specific adaptation for youth in foster care	Contraceptive use and consistency	12-months post intervention	High
Project IMAGE	Program for adolescents with a history of STIs and abuse	Health clinic	2, 3-4 hour sessions followed by 3-5, 1 hour support group sessions and 2 or more individual counseling sessions	African-American and Mexican-American adolescent girls ages 14-18 with a history of STIs and abuse	Sexually transmitted infections or HIV	12-months post intervention	Moderate

Evidence-Based TPP Programs are Implementation Ready and Eligible for Replication with OAH Funding

Program Model	Program Type	Implementation Setting	Implementation Length	Population	Outcomes	Duration of Outcomes	Evidence Review Rating
Project TALC	Program for parents living with HIV and their adolescent children	Community-based	24 sessions; 2-3 hours each; over 4-6 years	Parents living with HIV and their adolescent children	Pregnancy or birth	4 years after program start	High
Promoting Health Among Teens! Abstinence-Only Intervention	Abstinence education	In-school middle school After-school Community-based	8, 1-hour sessions	Youth ages 11-13	Sexual initiation or abstinence Recent sexual activity	Ranges, 3-24 months post intervention	High
Promoting Health Among Teens! Comprehensive Abstinence and Safer Sex Intervention	Sexuality education	In-school middle school After-school Community-based	12, 45-1-hour sessions	Youth ages 11-13	Number of sexual partners	Ranges, 3-12 months post intervention	High
Raising Healthy Children	Youth development	In-school	Multi-year, multi-session; includes program components for teachers and school staff, parents, and students	Schools grades K-6	Sexual initiation or abstinence Number of sexual partners Pregnancy or birth Sexually transmitted infections or HIV	Ranges from when participants were 18 years old to 27 years old	Moderate
Reducing the Risk	Sexuality education	In-school high school Community-based	16, 45-min lessons	Youth ages 13-18 in grades 8-12	Contraceptive use and consistency	18-months post intervention for girls only	Moderate
Respeto/Proteger	Program for Latino parenting teens	After school Community-based	6, 2-hour sessions	Young Latino parents with children at least 3 months of age	Contraceptive use and consistency	Ranges, 3-6 months post intervention	Moderate

Evidence-Based TPP Programs are Implementation Ready and Eligible for Replication with OAH Funding

Program Model	Program Type	Implementation Setting	Implementation Length	Population	Outcomes	Duration of Outcomes	Evidence Review Rating
<u>Rikers Health Advocacy Program (RHAP)</u>	Program for incarcerated youth	Correctional facility	4, 1-hour sessions	Incarcerated, inner-city adolescent males ages 16-19	Contraceptive use and consistency	10-months after program start	Moderate
<u>Safer Choices</u>	Sexuality education	In-school high school	11 sessions in 9 th grade, 10 sessions in 10 th grade	9 th and 10 th grade students	Contraceptive use and consistency	Two years after program start	High
<u>Safer Sex</u>	Clinic-based	Clinic-based	1, 30-50 minute session followed by 10-30 minute booster sessions at 1-month, 3-months, and 6-months	Sexually active female adolescents	Number of partners	6-months post intervention	Moderate
<u>Sexual Health and Adolescent Risk Prevention (SHARP) (formerly known as HIV Risk Reduction Among Detained Adolescents)</u>	Program for youth in juvenile detention	Juvenile detention facilities	1, 3-hour session followed by 1-hour group discussion of alcohol use and sexual activity	High-risk adolescents in juvenile detention facilities	Contraceptive use and consistency	12-months post intervention	Moderate
<u>SiHLE</u>	Sexuality education	Community-based	4 sessions with 6-month and 12-month follow-up sessions	African-American female teens ages 14-18	Contraceptive use and consistency Pregnancy or birth	12-months post intervention	High
<u>Sisters Saving Sisters</u>	Sexuality education	Community-based Clinic-based In schools	1, 4.5 hour session	Female adolescents ages 12-19	Number of sexual partners Contraceptive use and consistency Sexually transmitted infections or HIV	12-months post intervention	High

Evidence-Based TPP Programs are Implementation Ready and Eligible for Replication with OAH Funding							
Program Model	Program Type	Implementation Setting	Implementation Length	Population	Outcomes	Duration of Outcomes	Evidence Review Rating
STRIVE	Program for youth who have recently run away from home	Community-based	5, 90-120 minute sessions for the adolescent and at least one parent	Newly homeless youth ages 12-17	Number of sexual partners	12-months post intervention	Moderate
Teen Health Project	Sexuality education	Community-based Low-income housing developments	2, 3-hour workshops; 2, 90-120 min follow up session; 1, 90-min parent session; various community activities	Youth ages 12-17	Sexual initiation or abstinence	12-months post intervention	Moderate
Teen Outreach Program (TOP)	Youth development	In-school After-school Community-based	25 weekly session over a 9-month period and 20 hours of community service learning	Youth ages 12-19	Pregnancy or birth	Post-intervention	High
17 Days (formerly known as 'What Could You Do?')	Clinic-based	Health clinic	45-minute or longer video with 3-month and 6-month booster sessions	Female adolescents ages 14-18	Recent sexual activity Sexually transmitted infections or HIV	6-months post intervention	High

*This list is inclusive of only the evidence-based TPP programs eligible for replication at the time the FOA was published. Additional evidence-based TPP programs identified by the HHS TPP Evidence Review prior to the end of the planning and readiness period will also be eligible for replication under this FOA as long as they have been assessed by the HHS TPP Evidence Review as being implementation ready, meaning that the program has clearly defined curricula and components, necessary staff supports and training, and specified guidelines and tools for monitoring fidelity.