

We use cookies to enhance your experience on our website. By clicking 'continue' or by continuing to use our website, you are agreeing to our use of cookies. You can change your cookie settings at any time.

Continue
Find out more

OXFORD
ACADEMIC



Outbreak of Measles Among Persons With Prior Evidence of Immunity, New York City, 2011 ^{FREE}

Jennifer B. Rosen , Jennifer S. Rota, Carole J. Hickman, Sun B. Sowers, Sara Mercader, Paul A. Rota, William J. Bellini, Ada J. Huang, Margaret K. Doll, Jane R. Zucker, ... [Show more](#)

Clinical Infectious Diseases, Volume 58, Issue 9, 1 May 2014, Pages 1205–1210, <https://doi.org/10.1093/cid/ciu105>

Published: 27 February 2014 **Article history** ▼

A measles outbreak occurred in New York City. All cases had prior evidence of measles immunity. Symptoms were consistent with measles. Laboratory results indicated secondary immune responses. This report documents measles transmission from an individual with verified secondary vaccine failure.

Background. Measles was eliminated in the United States through high vaccination coverage and a public health system able to rapidly respond to measles. Measles may occur among vaccinated individuals, but secondary transmission from such individuals has not been documented.

Methods. Suspected patients and contacts exposed during a measles outbreak in New York City in 2011 were investigated. Medical histories and immunization records were obtained. Cases were confirmed by detection of measles-specific immunoglobulin M and/or RNA. Tests for measles immunoglobulin G (IgG), IgG avidity, measurement of measles neutralizing antibody titers, and genotyping were performed to characterize the cases.

Results. The index patient had 2 doses of measles-containing vaccine; of 88 contacts, 4 secondary patients were confirmed who had either 2 doses of measles-containing vaccine or a past positive measles IgG antibody. All patients had laboratory confirmation of measles infection, clinical symptoms consistent with measles, and high-avidity IgG antibody characteristic of a secondary immune response. Neutralizing antibody titers of secondary patients reached >80 000 mIU/mL 3–4 days after rash onset and that of the index was <500 mIU/mL 9 days after rash onset. No additional cases of measles occurred among 231 contacts of secondary patients.

Conclusions. This is the first report of measles transmission from a twice-vaccinated individual with documented secondary vaccine failure. The clinical presentation and laboratory data of the index patient were typical of measles in a naive individual. Secondary patients had robust anamnestic antibody responses. No tertiary cases occurred despite numerous contacts. This outbreak underscores the need for thorough epidemiologic and laboratory investigation of suspected cases of measles regardless of vaccination status.

Keywords: [measles](#), [outbreak](#), [immunity](#), [vaccine failure](#), [waning immunity](#)

Topic: [disease outbreaks](#), [immunity](#), [immunologic memory](#), [new york city](#), [vaccination](#), [measles](#), [infection after vaccination](#)

Issue Section: [Articles and Commentaries](#)

Before the introduction of measles vaccine, >90% of the US population contracted measles by age 15 years [1]. Following the introduction of measles vaccine in 1963, measles incidence declined rapidly [2]. In 1989, the Advisory Committee on Immunization Practices (ACIP) [skipped](#) recommended a second dose of measles vaccine, as combined measles–mumps–rubella (MMR) vaccine, for introduction into the routine immunization schedule [3]. Adherence to this recommendation, combined with sustained, high national MMR vaccination coverage, helped eliminate endemic measles transmission by 2000 [4, 5]. Measles remains endemic in many parts of the world, and international travelers with

[Article Navigation](#)

cause of measles following importation [6]. The ongoing risk of importations requires sustaining high levels of population immunity to maintain measles elimination in the United States.

The ACIP criteria for presumptive evidence of immunity to measles include documented age-appropriate receipt of live measles virus-containing vaccine, laboratory evidence of immunity, laboratory evidence of disease, or birth before 1957 [6, 7]. Although vaccination with 2 doses of MMR vaccine is highly effective and is a proxy for immunity to measles, cases of measles have occurred among persons despite receipt of 2 doses of MMR vaccine [8–13]. Persons with detectable but low levels of neutralizing antibody despite receipt of 2 doses of MMR vaccine are potentially susceptible to infection and disease [14–18]. When measles is introduced into a highly vaccinated population, there are fewer cases of measles; however, among the cases that occur, the relative proportion occurring in vaccinated individuals increases [19]. These cases generally occur in an outbreak or in a setting involving intense exposure to an unvaccinated person with measles, and often exhibit modified symptoms.

Although antibody levels are expected to decline over time, it is unclear what effect the absence of natural boosting (asymptomatic secondary immune response) by circulating virus may have in the future on the maintenance of overall population immunity, including the ability of vaccinated persons to transmit virus [20, 21]. Subsequent spread of disease has not been documented from measles patients with a verified secondary immune response [9, 12, 13].

We report on an outbreak of 5 cases of measles in New York City (NYC) in which a fully vaccinated index patient transmitted measles infection to 4 contacts with presumptive evidence of measles immunity.

METHODS

Patient and Contact Ascertainment and Investigation

The index patient was identified through routine surveillance at the NYC Department of Health and Mental Hygiene (DOHMH). Patients with measles were identified by mandatory healthcare provider reports and electronic laboratory notification of positive measles immunoglobulin M (IgM) and/or RNA results. Provider reports of immunization records and measles immunoglobulin G (IgG) titers done before the onset of illness were reviewed for all patients, if available; verbal reports were not accepted as documentation of immunity. Clinical information and lists of exposed contacts were obtained from a review of medical records and through patient interview. The 2009 Council of State and Territorial Epidemiologists case definition was used to classify confirmed cases [22]. A list of contacts was developed based on identifiable individuals at known exposure sites in NYC. Documented immunization records and measles IgG titers of identified contacts were reviewed and immunity to measles was determined based on ACIP criteria [7]. Contacts were informed about symptoms of measles and were instructed to contact the DOHMH if they developed measles symptoms. Follow-up with nonimmune contacts was conducted again at the end of the incubation period to assure that contacts remained asymptomatic. Exposures occurring outside of NYC were not included in this evaluation.

Laboratory Testing

Initial serological and virological testing was performed in several commercial and public health department laboratories. Subsequently, all specimens were sent to the MMR Laboratory at the Centers for Disease Control and Prevention (CDC) for confirmation. Only results from the MMR laboratory at CDC are presented.

Serum specimens were tested for measles-specific IgM antibodies using an IgM capture enzyme immunoassay (EIA) as previously described [23]. Measles-specific IgG was tested using an in-house indirect EIA assay incorporating the measles nucleoprotein as the antigen [23]. IgM/IgG index ratios were derived by dividing the net absorbance values measured for IgM by IgG [24]. The IgM/IgG ratio was compared among patients as a measure of primary vs secondary immune response to infection. Index ratios >1 suggested a primary immune response to measles, and ratios <1 were consistent with a secondary response [24].

Avidity of measles-specific IgG antibody was tested by modification of a commercial measles IgG EIA (Captia Measles IgG, Trinity Biotech, Jamestown, New York), as previously described [25]. Measles neutralizing antibody titers were measured using a plaque reduction neutralization (PRN) test performed as previously described [26–28]. Serum specimens were run in parallel with the World Health Organization (WHO) Second International Standard Anti-Measles serum (coded 66/202, supplied by National Institute for Biological Standards and Control, South Mimms, United Kingdom). According to the run validation parameters, a titer of 1:8 corresponded to 8 mIU/mL. Seropositivity was defined as PRN concentrations ≥ 8 mIU/mL and seroprotection ≥ 120 mIU/mL [28].

Reverse transcription polymerase chain reaction (RT-PCR) and genotyping were performed. RNA was extracted from nasopharyngeal swab specimens using the QIAamp Viral RNA Mini Kit (Qiagen, Gaithersburg, Maryland). Measles virus RNA was detected by using a real-time RT-

RESULTS

The index patient was a 22-year-old female resident of NYC with a past medical history only significant for mitral valve prolapse. She developed a generalized rash, cough, conjunctivitis, coryza, sore throat, and subjective fever and presented to an emergency room for medical care but was not hospitalized. She had documentation of receipt of MMR vaccination at 3 years and 4 years of age. There was no travel during the incubation period and no known sick contacts. However, the index patient worked at a theater frequented by tourists. Eighty-eight exposed contacts aged 20–65 years were identified in NYC during her infectious period, of whom 66 (75%) had documentation of immunity, 10 (11%) were not immune to measles at the time of exposure, and 12 (14%) had unknown immune status.

Subsequently, 4 additional patients were identified among contacts of the index patient (Table 1). Three of the secondary patients (cases 2, 4, and 5) were healthcare workers at a clinic where the index patient received care and were exposed to the index patient on her day of rash onset. The other secondary patient (case 3) was a coworker of the index patient and was exposed to the index patient 2 days prior to her rash onset. The secondary patients had no epidemiologic links to any other patient with measles in NYC. These secondary patients had a generalized rash with onset between days 12 and 16 after exposure to the index patient and ranged in age from 20 to 52 years (median, 30 years). Two of the secondary patients had 2 documented doses of MMR vaccine and 2 had prior positive measles IgG antibody results (Table 1). One patient (case 2) had a medical history of immunosuppression and presented with rash, fever, cough, coryza, and conjunctivitis (Table 1). All other secondary patients presented with either rash and fever (case 3) or rash, fever, and cough (cases 4 and 5). None were hospitalized and there were no complications (Table 1).

Table 1.

Medical History, Immunity History, and Clinical Presentation for Patients in a Measles Outbreak in New York City, 2011

| Case | Age, y | Prior Evidence of Immunity | | Medical History and Clinical Presentation | | | | | | | |
|----------------|--------|----------------------------|-------------------|--|-------------------|------------------|------------|-------|----------------|--------|------------------------|
| | | MMR, Year | Measles IgG, Year | Medical History | Rash ^a | Rash Duration, d | Fever | Cough | Conjunctivitis | Coryza | Other |
| 1 (index) | 22 | 1991; 1992 | ... | Mitral valve prolapse | Y | 8 | Subjective | Y | Y | Y | Sore throat |
| 2 | 25 | 1987 1990 | ... | Ulcerative colitis, immunosuppressive medication, cerebral palsy | Y | Unknown | 38.9°C | Y | Y | Y | Diarrhea; Koplik spots |
| 3 ^b | 20 | 1992 1996 | ... | None | Y | 4 | 38.9°C | N | N | N | Sore throat |
| 4 | 35 | ... | 2006 | None | Y | 3 | 38.3°C | Y | N | N | N |
| 5 ^c | 52 | ... | 1993 | Hypothyroid on Synthroid | Y | 5 | Subjective | Y | N | N | N |

Abbreviations: IgG, immunoglobulin G; MMR, measles-mumps-rubella.

^a Generalized.

^b Case was reported to the New York City Department of Health and Mental Hygiene (NYC DOHMH) but investigated by the jurisdiction of residence, Baltimore County Department of Health.

^c Case was reported to the NYC DOHMH but investigated by the jurisdiction of residence, Westchester County Department of Health.

An additional 231 contacts were identified as exposed to the secondary patients in NYC. Among these exposed contacts, 157 (68%) were considered to be immune to measles or received postexposure prophylaxis, 5 (2%) were not immune, and 69 (30%) had unknown immune status. No tertiary cases were identified among these contacts.

[Skip to Main Content](#)

[Article Navigation](#)

serum samples except 1 sample collected on day 0 from case 2, although the PRN titer from the same serum sample was 1367 mIU/mL and by day 4 IgG was positive and the PRN titer had increased by >100-fold. The ratio of IgM to IgG in the day 9 serum from the index patient was 9.7, whereas the other 4 patients had ratios <1.0 (Table 2). Measles RT-PCR was positive from nasopharyngeal specimens from 4 of the 5 patients, 2 of which were sequenced and identified as genotype D4 (Table 2).

Table 2.

Laboratory Results From Serologic and Virologic Testing of Patients in a Measles Outbreak in New York City, 2011

| Case | Serum Sample No.: Days From Rash Onset to Collection | IgM Result | IgG Result | IgM/IgG Index Ratio ^a | IgG Avidity (etAI) ^b | PRN Titer, mIU/mL | RT-PCR | Genotype |
|----------------|--|------------|------------|----------------------------------|---------------------------------|-------------------|----------|----------|
| 1 | Serum 1: 2 d | Positive | Positive | ... | Intermediate (63%) | 81 | Positive | D4 |
| | Serum 2: 9 d | Positive | Positive | 9.7 | High (100%) | 402 | | |
| 2 | Serum 1: 0 d | Negative | Negative | ... | Not done | 1367 | Positive | D4 |
| | Serum 2: 4 d | Positive | Positive | ... | High (83%) | 150 219 | | |
| | Serum 3: 11 d | Positive | Positive | 0.3 | High (79%) | 175 563 | | |
| 3 ^c | Serum 1: 3 d | Positive | Positive | ... | High (82%) | 87 155 | Positive | Not done |
| | Serum 2: 6 d | Positive | Positive | ... | High (70%) | 221 291 | | |
| | Serum 3: 11 d | Positive | Positive | 0.2 | High (73%) | 168 036 | | |
| 4 | Serum 1: 3 d | Positive | Positive | ... | Intermediate (62%) | 107 712 | Negative | Not done |
| | Serum 2: 10 d | Positive | Positive | 0.24 | High (70%) | 94 860 | | |
| 5 ^d | Serum 1: 3 d | Negative | Positive | ... | High (92%) | 94 860 | Positive | Not done |
| | Serum 2: 7 d | Negative | Positive | 0.02 | High (97%) | 171 632 | | |

Abbreviations: etAI, end-titer avidity index; IgG, immunoglobulin G; IgM, immunoglobulin M; PRN, plaque reduction neutralization; RT-PCR, reverse transcription polymerase chain reaction.

^a IgM/IgG index ratio calculated on latest serum available for each patient (7–11 days); index ratios >1 suggest a primary immune response to measles and ratios <1 suggest a secondary response.

^b Avidity classified as low (etAI ≤30%), intermediate (etAI 31%–70%), and high (etAI ≥70%); intermediate and high avidity suggest past immunologic experience with measles through vaccination or natural measles infection.

^c Case reported to the New York City Department of Health and Mental Hygiene (NYC DOHMH) but investigated by the Baltimore County Department of Health.

^d Case reported to the NYC DOHMH but investigated by jurisdiction of residence, Westchester County Department of Health.

The initial serum samples collected from the index patient and case 4 had IgG avidity measured in the intermediate range (end-titer avidity index: 63% and 62%, respectively). However, all of the patients had high-avidity IgG in follow-up serum samples (Table 2). The PRN titer obtained from the index patient was 81 mIU/mL in serum collected 2 days after rash and 402 mIU/mL in the follow-up serum collected 7 days later. The 4 secondary patients had PRN titers of >80 000 mIU/mL in serum collected at ≥3 days after rash onset.

DISCUSSION

An unusual outbreak of measles was investigated in which all of the patients had either 2 documented doses of MMR vaccine or a positive result recorded for measles IgG antibody in the past. The index patient had 2 documented doses of MMR vaccine before infection and subsequently transmitted disease to 4 contacts. Although other outbreaks have been reported in which persons with a history of MMR vaccination were confirmed with measles, this is the first report in which a person with a verified secondary vaccine failure despite receipt of two doses of MMR was demonstrated to be capable of transmitting disease to other individuals [8–13, 33].

[Skip to Main Content](#)

[Article Navigation](#)

was typical of a primary response, whereas those of the 4 secondary patients were consistent with a secondary immune response [14, 24]. Although the detection of IgM among measles patients had long been presumed to be a characteristic of the first exposure to measles antigen (ie, primary immune response), it has since been recognized that IgM can be elicited by persons with either a symptomatic or asymptomatic secondary immune response, albeit at lower IgM to IgG ratios [14, 24]. Despite high-avidity IgG, the index patient did not develop the typical high neutralizing antibody titers that have been observed among previously immunized patients [12, 13]. Previous studies have noted that extremely high PRN titers in acute-phase serum from vaccinated persons with suspected measles infection might serve as a biomarker for patients with a secondary immune response [12, 13]. The 4 secondary patients all had an early and robust antibody response within a few days after rash onset, with PRN titers 6–60 times higher than those observed after primary infection with natural disease or following measles vaccination [12, 13]. Despite >200 exposures identified through investigations following notification of the 4 secondary patients, no additional cases were detected. This is in agreement with other published investigations describing a lack of transmission by documented cases of secondary vaccine failure [12, 13]. In each of the secondary patients, neutralizing antibody may have waned sufficiently to allow symptomatic infection, but the anamnestic response upon reexposure to measles generated a rapid and robust memory response that may have reduced their infectious period. However, in the setting of high population-level immunity, it is challenging to evaluate transmissibility. In contrast, the index patient's lower neutralizing antibody titer after infection provides a biologically plausible explanation for her ability to transmit virus.

Previous outbreaks in which symptomatic patients with measles who were later confirmed to have a secondary immune response generally involved patients with modified clinical presentations that could easily have been misdiagnosed in the absence of another confirmed case of measles [12, 13]. However, based on the clinical information available, the index patient and 3 of the 4 secondary patients had typical clinical presentations of a generalized rash with duration of 3 or more days with fever, and either cough, conjunctivitis, or coryza.

An understanding of the duration of measles immunity is important for ensuring continued success with global measles elimination. Demonstration of waning immunity requires measurement of neutralizing antibody titers before revaccination or infection and at time points subsequent to the antigenic stimulation. Detection of IgG is a proxy for immunity, not an absolute correlate of protection from disease. Similarly, the inability to detect measles-specific IgG should not be construed as a lack of immunity in persons who have been previously vaccinated, as cellular immunity and antibody functionality play important roles in protection [17, 34–37]. Neutralizing antibody titers have the closest correlation with immunity to measles, but assays that measure measles neutralizing antibody are not widely available, and tests of cellular immunity are challenging to perform [36, 38].

With the achievement of measles elimination, boosting of immunity from exposure to wild-type measles virus is uncommon [20]. It is unknown if boosting of immunity among vaccinated individuals by exposure to circulating virus had previously played an important role in maintaining protective levels of antibody, raising questions about the duration of population immunity to measles. In one report of schoolchildren in a postelimination environment, measles neutralizing antibodies persisted for 10 years after receipt of a second dose of MMR. Although no seronegative results were detected after 10 years, titers did decline over time, with 4.7% (18/382) of the children considered potentially susceptible to infection, given PRN titers in the range of 8–120 mIU/mL; however, 72% of those with low titers after 10 years had been in the lowest quartile of titers prior to the second MMR dose [17].

There are limitations to this evaluation. Although provider documentation of MMR vaccination was obtained, it is not possible to know about the quality of the vaccine received. Inappropriate storage conditions could alter the effectiveness of vaccine. It is possible that the patients never responded adequately or only achieved minimal titers following vaccination; however, the laboratory results of high-avidity IgG antibody demonstrate that all the patients had responded previously to measles virus and are not primary vaccine failures.

As we move forward on global efforts to eliminate measles, it remains critical that we maintain high population immunity and vigilance for disease. International importations of measles continue to occur in the United States [6]. The current 2-dose MMR vaccination strategy has successfully maintained measles elimination in the United States for nearly 20 years, despite continued importations. Now that the United States has been free of endemic measles for more than a decade and natural boosting of infection is uncommon, it will be important to better understand the duration of immunity [20, 39, 40]. However, this outbreak probably represents a series of rare events, and waning immunity among previously vaccinated persons is unlikely to threaten the ability to sustain measles elimination.

Surveillance also plays a vital role in monitoring the status and duration of population immunity by identifying instances of disease in individuals with prior immunity and conducting investigations of their exposed contacts. Although 3 of the 4 secondary patients described in this report had typical clinical presentations, it is important to note that previous outbreaks in which measles patients with a secondary immune response were identified generally involved patients with modified clinical presentations that could easily have been misdiagnosed in the absence of a thorough investigation [12, 13]. A single episode of transmission from an individual with prior evidence of immunity does not justify a change in current measles control and elimination strategies; however, this case clearly underscores the need to maintain sensitive surveillance activities.

[Skip to Main Content](#)

Notes

[Article Navigation](#)

Sihly, of the New York City Department of Health and Mental Hygiene, David Blythe of the Maryland Department of Health and Mental Hygiene, and Laurie Smalley and India Harris of the Westchester County Department of Health for conducting case and contact investigations, and Laura Walls, Nobia Williams, Rebecca McNall, Elena Lopareva, and Luis Lowe from the Centers for Disease Control and Prevention for laboratory support.

Potential conflicts of interest. All authors: No reported conflicts.

All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

- 1 Langmuir A Medical importance of measles *Am J Dis Child* 1962;103:54–6
- 2 Orenstein WA, Papania MJ, Wharton ME. Measles elimination in the United States. *J Infect Dis* 2004;189 suppl 1 S1–3
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 3 Centers for Disease Control and Prevention. Measles, mumps, and rubella—vaccine use and strategies for elimination of measles, rubella, and congenital rubella syndrome and control of mumps: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Morb Mortal Wkly Rep* 1998;47:1–57
[PubMed](#)
- 4 Centers for Disease Control and Prevention. National, state, and local area vaccination coverage among children aged 19–35 months—United States, 2011. *MMWR Morb Mortal Wkly Rep* 2012;61:689–96
[PubMed](#)
- 5 Katz SH, Hinman A. Summary and conclusions: measles elimination meeting 16–17 March 2000. *J Infect Dis* 2004;189 suppl 1 S43–7
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 6 Centers for Disease Control and Prevention. Measles—United States, 2011. *MMWR Morb Mortal Wkly Rep* 2012;61:253–7
[PubMed](#)
- 7 Centers for Disease Control and Prevention. Prevention of measles, rubella, congenital rubella syndrome, and mumps, 2013. *MMWR Morb Mortal Wkly Rep* 2013;62:1–34
[PubMed](#)
- 8 Ammari LB, Bell LH, Hodinka R. Secondary measles vaccine failure in healthcare workers exposed to infected patients. *Infect Control Hosp Epidemiol* 1993;14:81–6
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 9 Atrasheuskaya AK, Kulak MN, Neverov AR, Rubin S, Ignatyev G. Measles cases in highly vaccinated population of Novosibirsk, Russia, 2000–2005. *Vaccine* 2008;26:2111–8
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 10 Coleman K, Markey P. Measles transmission in immunized and partially immunized air travellers. *Epidemiol Infect* 2010;138:1012–5
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 11 Edmonson MA, Adonis DM, McPherson JB, Berg JC, Circo S, David JM. Mild measles and secondary vaccine failure during a sustained outbreak in a highly vaccinated population. *JAMA* 1990;263:2467–71
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 12 Hickman CJ, Hyde TB, Sowers SB, et al. Laboratory characterization of measles virus infection in previously vaccinated and unvaccinated individuals. *J Infect Dis* 2011;204 suppl 1 S549–58
[Google Scholar](#) [Crossref](#) [PubMed](#)

[Skip to Main Content](#)

Article Navigation

[Google Scholar](#) [Crossref](#) [PubMed](#)

- 14 Chen RTMarkowitz LEAlbrecht Pet al. Measles antibody: reevaluation of protective titers*J Infect Dis* 19901621036 42
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 15 Haralambieva IHOvsyannikova IGO'Byrne MPankratz VSJacobson RMPoland GAA large observational study to concurrently assess persistence of measles specific B-cell and T-cell immunity in individuals following two doses of MMR vaccine*Vaccine* 2011294485 91
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 16 Haralambieva IHOvsyannikova IGPankratz VSKennedy RBJacobson RMPoland GAThe genetic basis for interindividual immune response variation to measles vaccine: new understanding and new vaccine approaches*Expert Rev Vaccines* 20131257 70
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 17 LeBaron CWBeeler JASullivan BJet al. Persistence of measles antibodies after 2 doses of measles vaccine in a postelimination environment*Arch Pediatr Adolesc Med* 2007161294 301
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 18 Poland GJacobson RThampy Aet al. Measles reimmunization in children seronegative after initial immunization*JAMA* 19972771156 8
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 19 Centers for Disease Control and PreventionMeasles vaccine efficacy—United States*MMWR Morb Mortal Wkly Rep* 198029470 2
- 20 Mossong JMuller CModelling measles re-emergence as a result of waning of immunity in vaccinated populations*Vaccine* 2003214597 603
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 21 Pebody RGGay NJHesketh LMet al. Immunogenicity of second dose measles-mumps-rubella (MMR) vaccine and implications for serosurveillance*Vaccine* 2002201134 40
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 22 CSTE position statement 09-ID-48: public health reporting and national notification for measlesAvailable at: <http://c.ymcdn.com/sites/www.cste.org/resource/resmgr/PS/09-ID-48.pdf> Accessed 8 March 2014
- 23 Hummel KBERdman DDHeath JBellini WJBaculovirus expression of the nucleoprotein gene of measles virus and utility of the recombinant protein in diagnostic enzyme immunoassays*J Clin Microbiol* 1992302874 80
[Google Scholar](#) [PubMed](#)
- 24 Erdman DDHeath JLWatson JCMarkowitz LEBellini WJImmunoglobulin M antibody response to measles virus following primary and secondary vaccination and natural virus infection*J Med Virol* 19934144 8
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 25 Mercader SGarcia PBellini WMeasles IgG avidity assay: use in classification of measles vaccine failure cases in elimination settings*Clin Vaccine Immunol* 2012191810 7
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 26 Albrecht PHerrmann KBurns GRRole of virus strain in conventional and enhanced measles plaque neutralization test*J Virol Methods* 19813251 60
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 27 Lennette EHSchmidt N*Diagnostic procedures for viral and rickettsial infections* 19795th edWashington, DCAmerican Public Health Association
- 28 Cohen BAudet SAndrews NBeeler JPlaque reduction neutralization test for measles antibodies: description of a standardised laboratory method for use in immunogenicity studies of aerosol vaccination*Vaccine* 20072659 66
[Google Scholar](#) [Crossref](#) [PubMed](#)
- [Skip to Main Content](#)
- 29 Hummel KBLowe LBellini WJRota PADevelopment of quantitative gene-specific real-time RT-PCR assays for the detection of measles virus in clinical specimens*J Virol Methods* 2006132166 73

Article Navigation

- 30 Rota PABrown KEHubschen JMet al. Improving global virologic surveillance for measles and rubella*J Infect Dis* 2011204 suppl S1 506 13
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 31 Rota PBrown KMankertz Aet al. Global distribution of measles genotypes and measles molecular epidemiology*J Infect Dis* 2011204 suppl 1 S514 23
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 32 World Health OrganizationMeasles virus nomenclature update: 2012*Wkly Epidemiol Rec* 20128773 81
[PubMed](#)
- 33 Yeung LFLurie PDayan Get al. A limited outbreak in a highly vaccinated US boarding school*Pediatrics* 20051161287 91
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 34 Gans HYasukawa LAllderson Aet al. Humoral and cell-mediated immune responses to an early 2-dose measles vaccination regimen in the United States*J Infect Dis* 200419083 90
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 35 Gans HYasukawa LRinki Met al. Immune responses to measles and mumps vaccination of infants at 6, 9, and 12 months*J Infect Dis* 2001184817 26
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 36 Strebel PPapania MFiebelkorn AHalsey NPlotkin SOrenstein WOffit PMeasles vaccine*Vaccines* 2012PhiladelphiaElsevier352 87
- 37 Ward BBoulianne NRatnam SGulut M-CCouillard MDeSerres GCellular immunity in measles vaccine failure: demonstration of measles antigen-specific lymphoproliferative responses despite limited serum production after revaccination*J Infect Dis* 19951721591 5
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 38 Pichichero MBooster vaccinations: can immunologic memory outpace disease pathogenesis?*Pediatrics* 20091241633 41
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 39 Mossong JNokes DEdmunds Wcox MRatnam SMuller CModeling the impact of subclinical measles transmission in vaccinated populations with waning immunity*Am J Epidemiol* 19991501238 49
[Google Scholar](#) [Crossref](#) [PubMed](#)
- 40 van den Hof SBerbers Gde Melker HConyn-van Spaendonck MMarin MSero-epidemiology of measles antibodies in the Netherlands, a cross-sectional study in a national sample and in communities with low vaccine coverage*Vaccine* 199918931 40
[Google Scholar](#) [Crossref](#) [PubMed](#)

© The Author 2014. Published by Oxford University Press on behalf of the Infectious Diseases Society of America. All rights reserved. For Permissions, please e-mail: journals.permissions@oup.com.

Comments

0 Comments

[Add comment](#)

[Skip to Main Content](#)

[Article Navigation](#)[View Metrics](#)

Email alerts

[New issue alert](#)[Advance article alerts](#)[Article activity alert](#)

Receive exclusive offers and updates
from Oxford Academic

More on this topic

Immunity to Measles, Mumps, and Rubella in US Children With Perinatal HIV Infection or Perinatal HIV Exposure Without Infection

Lessons from the West Nile Viral Encephalitis Outbreak in New York City, 1999: Implications for Bioterrorism Preparedness

Clinical Characteristics of Measles in Previously Vaccinated and Unvaccinated Patients in California

Birth Cohort Studies Assessing Norovirus Infection and Immunity in Young Children: A Review

Related articles in

[Web of Science](#)

[Google Scholar](#)

Related articles in PubMed

Acute otitis media in infants younger than two months of age: Epidemiologic and microbiologic characteristics in the era of pneumococcal conjugate vaccines.

[Skip to Main Content](#)

Declining prevalence of pediatric sudden deafness during the past two decades.

[Article Navigation](#)

Association between urban green space and the risk of cardiovascular disease: A longitudinal study in seven Korean metropolitan areas.

Citing articles via

Web of Science (39)

Google Scholar

CrossRef

Latest | **Most Read** | Most Cited

Inherited Interleukin 2–Inducible T-Cell Kinase Deficiency in Siblings With Epidermodysplasia Verruciformis and Hodgkin Lymphoma

Seroprotection at different levels of the health care system after routine vaccination with DTPw-HepB-Hib in Lao PDR

Risk of ectopic pregnancy and tubal infertility following gonorrhoea and chlamydia infections

Indirect Effects of 10-Valent Pneumococcal Conjugate Vaccine Against Adult Pneumococcal Pneumonia in Rural Western Kenya

Persistent Low-Level Viremia while on Antiretroviral Therapy is an Independent Risk Factor for Virologic Failure



[Skip to Main](#)

[About Clinical Infectious Diseases](#)[Editorial Board](#)[Author Guidelines](#)[Facebook](#)[Twitter](#)[Purchase](#)[Recommend to your Library](#)[Advertising and Corporate Services](#)[Journals Career Network](#)

Online ISSN 1537-6591

Print ISSN 1058-4838

Copyright © 2019 Infectious Diseases Society of America

[About Us](#)[Contact Us](#)[Careers](#)[Help](#)[Access & Purchase](#)[Rights & Permissions](#)[Open Access](#)

Resources

[Authors](#)[Librarians](#)[Societies](#)[Sponsors & Advertisers](#)[Press & Media](#)[Agents](#)

Connect

[Join Our Mailing List](#)[OUPblog](#)[Twitter](#)[Facebook](#)[YouTube](#)[Tumblr](#)

Explore

[Shop OUP Academic](#)[Oxford Dictionaries](#)[Oxford Index](#)[Epigeum](#)[OUP Worldwide](#)[University of Oxford](#)

Oxford University Press is a department of the University of Oxford. It furthers the University's objective of excellence in research, scholarship, and education by publishing worldwide

OXFORD
UNIVERSITY PRESS

Copyright © 2019 Oxford University Press

[Cookie Policy](#)

[Privacy Policy](#)

[Legal Notice](#)

[Site Map](#)

[Accessibility](#)

[Get Adobe Reader](#)