

Health Effects of Microwave Radio Exposures



Paul Dart MD FCA

The current FCC Limits for Microwave RF Exposure were published in 1999.

These guidelines are only designed to protect the public against the thermal effects of microwave RF.

The FCC has explicitly stated that they do not make any regulations or assurances whatsoever regarding the “nonthermal” biological effects of lower level microwave RF exposures (other physiologic effect besides heat damage).

Many statements from industry spokesmen state that “not enough is known” about these exposures to identify risk, or that there is “insufficient” or “incomplete” evidence regarding such risks, or that there is “no scientific consensus” on this risk.

This implies that there isn’t much scientific information on this subject. But actually, there is a great deal of research documenting adverse biological effects from low level RF exposures.

The FCC guidelines only address the thermal effects of RF exposure.



The current FCC Limits for Microwave RF Exposure were published in 1999. These guidelines are only designed to protect the public against the **thermal effects** of microwave RF.

The FCC has explicitly stated that they do not make any regulations or assurances whatsoever regarding the **“nonthermal” biological effects** of lower level microwave RF exposures (other physiologic effect besides heat damage).

Many statements from industry spokesmen state that “not enough is known” about these biological effects to identify risk, or that there is “insufficient” or “incomplete” evidence regarding such risks, or that there is “no scientific consensus” on this risk.

This implies that there isn’t much scientific information on this subject. But actually, there is a great deal of research documenting adverse biological effects from low level RF exposures.

Acute Symptoms from RF Exposure:

“EHS”



In this presentation, we're going to take a look at the current scientific evidence for adverse biological effects of RF exposures.

First, we'll look at the evidence that RF exposures can produce **acute symptoms** in many individuals, a problem called **Electrohypersensitivity Syndrome** or “EHS”.

Then we'll look at the evidence that RF exposures **alter hormone physiology** and **increase oxidative stress** in living systems.

Then we'll review the evidence that such alterations in physiology can **damage DNA**, increase the incidence of some forms of **cancer**, and decrease **fertility** in animals and in human beings.

U.S. Embassy, Moscow (1953-1978)



Acute symptoms provoked by microwave radiation were first described by Russian medical researchers in the 1950's. They described a constellation of symptoms including headache, ocular dysfunction, fatigue, dizziness, sleep disorders, dermatographism, cardiovascular abnormalities, depression, irritability, and memory impairment.

In the years **between 1953 and 1978** the Russian government harrassed the U.S. Embassy in Moscow by targeting it with radiation from a microwave transmitter positioned on the roof of a nearby building.

Exposed embassy staff experienced a statistically significant excess of several problems, including: depression, irritability, difficulty in concentrating, memory loss, ear problems, skin problems, vascular problems, and other health problems. Symptom incidence increased significantly with accrued years of exposure. Exposure levels inside the building were in the order of **2 to 28 $\mu\text{W}/\text{cm}^2$ (FCC Guidelines: 600 $\mu\text{W}/\text{cm}^2$)**

U.S. Embassy, Moscow (1953-1978)

50272-101					
REPORT DOCUMENTATION PAGE	1. REPORT NO.	2.	3. Document Accession No. PB288163		
4. Title and Subtitle Evaluation of Health Status of Foreign Service and other Employees from Selected Eastern European Posts			5. Report Date July 31, 1978		
7. Author(s) Abraham M. Lilienfeld, M.D. Principal Investigator			6. Performing Organization Rept. No.		
9. Performing Organization Name and Address Department of Epidemiology School of Hygiene and Public Health The Johns Hopkins University			10. Project/Task/Work Unit No.		
12. Sponsoring Organization Name and Address Medical Director Office of Medical Services Department of State Washington, D C			11. Contract(C) or Grant(G) No. (C) 6025 619073 (G)		
13. Type of Report & Period Covered Final 1953 - 1976			14.		
15. Supplementary Notes Released publicly November 20 simultaneously by Department of State and The Johns Hopkins University.					
16. Abstract (Limit: 200 words) This is a biostatistical study of 1827 Department of State employees and their dependents at the Moscow Embassy and 2561 employees and their dependents from other Eastern European Embassies. Health records, health questionnaires and death certificates were the basic information sources. The study is the impact of the Moscow environment including microwave exposure on the health status and mortality of the employees. It was concluded that personnel working at the American Embassy in Moscow from 1953 to 1976 suffered no ill effects from the microwaves beamed at the Embassy.					

Concern about health effects among Embassy personnel led to a detailed study by A.M. Lilienfeld, an epidemiologist at Johns Hopkins University. The abnormalities found in this study were an embarrassment to the U.S. government, since the levels of exposure experienced by embassy staff inside the building were in the order of 2 to 28 microwatts/cm², a level dramatically below the described U.S. safety standards for microwave exposure. It appears that the conclusions of the study were altered to soft-pedal any abnormal findings.

Lilienfeld AM LGM, Cauthen J, Tonascia S, Tonascia J. Evaluation of health status of foreign service and other employees from selected eastern European embassies. *Foreign Service Health Status Study, Final Report*; Contract No. 6025-619037 (NTIS publication P8-288 163/9) (1979); 1-447.

Liakouris AG. Radiofrequency (RF) Sickness in the Lilienfeld Study: An Effect of Modulated Microwaves? *Archives of Environmental Health* (1998); 53(3):236-238.

Goldsmith JR. Where the trail leads. Ethical problems arising when the trail of professional work leads to evidence of a cover-up of serious risk and mis-representation of scientific judgement concerning human exposures to radar. *Eubios Journal of Asian and International Bioethics* (1995b); 5(4):92-94.

Cherry N. Evidence of Health Effects of Electromagnetic Radiation, To the Australian Senate Inquiry into Electromagnetic Radiation (2000): 1-84. http://www.neil-cherry.com/documents/90_m1_EMR_Australian_Senate_Evidence_8-9-2000.pdf

Norway (1998)



From:

Mild, K.H., Oftedal, G., Sandstrom, M., Wilen, J., Tynes, T., Haugsdal, B. and Hauger E., 1998: "Comparison of symptoms by users of analogue and digital mobile phones - A Swedish- Norwegian epidemiological study". National Institute for Working Life, 1998:23, Umea, Sweden, 84pp.

Norway (1998)




Figure 11: Prevalence of symptoms for Norwegian mobile phone users, mainly analogue, with various categories of length of calling time per day, from Mild et al. (1998).


In:

Cherry N. EMF/EMR Reduces Melatonin in Animals and People. (2002):1-14. <http://www.neilcherry.com/documents.php>

From:

Mild, K.H., Oftedal, G., Sandstrom, M., Wilen, J., Tynes, T., Haugsdal, B. and Hauger E., 1998: "Comparison of symptoms by users of analogue and digital mobile phones - A Swedish- Norwegian epidemiological study". National Institute for Working Life, 1998:23, Umea, Sweden, 84pp.

La Ñora, Spain (2001)



Town of 1900 inhabitants, with GSM cell phone tower.

Questionnaire distributed, 5% of inhabitants responded. The questionnaire was composed of 25 different items mainly concerning health information about the respondents.

The respondents scored and marked from 0 to 3 the presence of the suffered health dysfunction: 0 never, 1 sometimes, 2 often, 3 very often.


Power density of signal in bedrooms of respondents was measured.

Area A (< 150 meters from tower) = average power density $0.11 \mu\text{W}/\text{cm}^2$.

Area B (> 250 meters from tower) = average power density $0.01 \mu\text{W}/\text{cm}^2$.

Navarro E, Segura J, Portolés M, Gómez-Perretta C. The Microwave Syndrome: A Preliminary Study in Spain. *Electromagn Biol Med* (2003); 22(2-3):161-169.

La Ñora, Spain (2001)



Symptom score (0 – 3) vs Average Bedroom Exposure Levels to Microwave RF


FCC Guidelines: 600 - 1000 $\mu\text{W}/\text{cm}^2$

Navarro E, Segura J, Portolés M, Gómez-Perretta C. The Microwave Syndrome: A Preliminary Study in Spain. *Electromagn Biol Med* (2003); 22(2-3):161-169.

Abstract

A health survey was carried out in Murcia, Spain, in the vicinity of a Cellular Phone Base Station working in DCS-1800 MHz. This survey contained health items related to “microwave sickness” or “RF syndrome.” The microwave power density was measured at the respondents' homes. Statistical analysis showed significant correlation between the declared severity of the symptoms and the measured power density. The separation of respondents into two different exposure groups also showed an increase of the declared severity in the group with the higher exposure.

La Ñora, Spain (2001)



Symptom score (0 – 3) vs Average Bedroom Exposure Levels to Microwave RF

Based on the data of this study the advice would be to strive for levels not higher than 0.02 V/m for the sum total, which is equal to a power density of 0.0001 $\mu\text{W}/\text{cm}^2$ or 1 $\mu\text{W}/\text{m}^2$, which is the indoor exposure value for GSM base stations proposed on empirical evidence by the Public Health Office of the Government of Salzburg in 2002.

Oberfeld G, Navarro E, Portoles M, Maestu C, Gomez-Perretta C. The Microwave Syndrome -- Further Aspects of a Spanish Study. (2004):1-8. http://www.powerwatch.org.uk/pdfs/20040809_kos.pdf

France (2002)



Questionnaire re multiple symptoms that have been described for “microwave syndrome”.


n = 530

Evaluated incidence of symptoms as a function of residential proximity in meters to a cell phone tower.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49.

Fatigue



* p < 0.05 in comparison to residence > 300 meters or not exposed.

X axis = responses grouped by residential proximity to cell phone tower (in meters).


Y axis = percentage in exposure category answering "Often" or "Very Often" to a symptom query, in reference to total number of respondents living at that distance who answered "Never" to that symptom query.

* = statistically significant for this sample size.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49

Irritability



* p < 0.05 in comparison to residence > 300 meters or not exposed.

X axis = responses grouped by residential proximity to cell phone tower (in meters).


Y axis = percentage in exposure category answering "Often" or "Very Often" to a symptom query, in reference to total number of respondents living at that distance who answered "Never" to that symptom query.

* = statistically significant for this sample size.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49

Headache



* $p < 0.05$ in comparison to residence > 300 meters or not exposed.

X axis = responses grouped by residential proximity to cell phone tower (in meters).


Y axis = percentage in exposure category answering "Often" or "Very Often" to a symptom query, in reference to total number of respondents living at that distance who answered "Never" to that symptom query.

* = statistically significant for this sample size.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49

Sleep Disruption



* $p < 0.05$ in comparison to residence > 300 meters or not exposed.

X axis = responses grouped by residential proximity to cell phone tower (in meters).


Y axis = percentage in exposure category answering "Often" or "Very Often" to a symptom query, in reference to total number of respondents living at that distance who answered "Never" to that symptom query.

* = statistically significant for this sample size.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49

Depression



* p < 0.05 in comparison to residence > 300 meters or not exposed.

X axis = responses grouped by residential proximity to cell phone tower (in meters).


Y axis = percentage in exposure category answering "Often" or "Very Often" to a symptom query, in reference to total number of respondents living at that distance who answered "Never" to that symptom query.

* = statistically significant for this sample size.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49

Difficulty in Concentration



* $p < 0.05$ in comparison to residence > 300 meters or not exposed.

X axis = responses grouped by residential proximity to cell phone tower (in meters).


Y axis = percentage in exposure category answering "Often" or "Very Often" to a symptom query, in reference to total number of respondents living at that distance who answered "Never" to that symptom query.

* = statistically significant for this sample size.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49

Memory Loss



* $p < 0.05$ in comparison to residence > 300 meters or not exposed.

X axis = responses grouped by residential proximity to cell phone tower (in meters).


Y axis = percentage in exposure category answering "Often" or "Very Often" to a symptom query, in reference to total number of respondents living at that distance who answered "Never" to that symptom query.

* = statistically significant for this sample size.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49

Skin Problems



* $p < 0.05$ in comparison to residence > 300 meters or not exposed.

X axis = responses grouped by residential proximity to cell phone tower (in meters).


Y axis = percentage in exposure category answering “Often” or “Very Often” to a symptom query, in reference to total number of respondents living at that distance who answered “Never” to that symptom query.

* = statistically significant for this sample size.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49

Visual Disruption



* $p < 0.05$ in comparison to residence > 300 meters or not exposed.

X axis = responses grouped by residential proximity to cell phone tower (in meters).


Y axis = percentage in exposure category answering "Often" or "Very Often" to a symptom query, in reference to total number of respondents living at that distance who answered "Never" to that symptom query.

* = statistically significant for this sample size.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49

Hearing Disruption



* $p < 0.05$ in comparison to residence > 300 meters or not exposed.

X axis = responses grouped by residential proximity to cell phone tower (in meters).


Y axis = percentage in exposure category answering “Often” or “Very Often” to a symptom query, in reference to total number of respondents living at that distance who answered “Never” to that symptom query.

* = statistically significant for this sample size.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49

Dizziness



* $p < 0.05$ in comparison to residence > 300 meters or not exposed.

X axis = responses grouped by residential proximity to cell phone tower (in meters).


Y axis = percentage in exposure category answering "Often" or "Very Often" to a symptom query, in reference to total number of respondents living at that distance who answered "Never" to that symptom query.

* = statistically significant for this sample size.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49

Movement Difficulties



* $p < 0.05$ in comparison to residence > 300 meters or not exposed.

X axis = responses grouped by residential proximity to cell phone tower (in meters).


Y axis = percentage in exposure category answering "Often" or "Very Often" to a symptom query, in reference to total number of respondents living at that distance who answered "Never" to that symptom query.

* = statistically significant for this sample size.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49

Cardiovascular Problems



* $p < 0.05$ in comparison to residence > 300 meters or not exposed.

X axis = responses grouped by residential proximity to cell phone tower (in meters).

Y axis = percentage in exposure category answering “Often” or “Very Often” to a symptom query, in reference to total number of respondents living at that distance who answered “Never” to that symptom query.

* = statistically significant for this sample size.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49

Elderly people are more vulnerable

Table 3. Influence of age on the percentages of complaints

Symptoms	≤ 20 years		21–40 years		41–60 years		> 60 years	
	Distances of subjects from antennas (in meters)							
	≤ 300	> 300	≤ 300	> 300	≤ 300	> 300	≤ 300	> 300
Fatigue	56.7	62.5	82.4*	25	81.4*	57.8	73.3*	40
Irritability	16.2	11.1	46.2	18.2	50.5	35.3	52.1*	21
Headaches	42.4	26.3	57.6*	18.2	52*	13.3	49.5*	10
Nausea	2	0	12.9	0	9.9	0	15.6	15.7
Loss of appetite	13.3	8.8	12.7	0	11.8	0	15.9	15
Sleep disturbances	26.1	14.8	53*	12.5	73.9	52.6	68.5*	44.4
Depressive tendencies	10.2	5.7	14	5.8	36	20	41.7	27.7
Feeling of discomfort	4.4	2.9	26.3	6	41.6	16.6	45*	19
Difficulties in concentration	30.3	40	42.1	18.7	45.8	36.8	53.3*	20
Memory loss	7.5	8	21.8	6.6	43	40	64	36.8
Skin problems	16.6	9.3	24.2	6.6	18.3	0	20.4	5.2
Visual disturbances	16.3	12.5	14.7	12.5	26.6	26.3	36.8	17.6
Hearing disturbances	9.4	5.1	15.4	0	29.8	21.7	43.8	31.5
Dizziness	6.2	5.2	3.2	6.6	15.4	4.5	39.3*	9.5
Movement difficulties	0	2.3	0	0	3.5	4	21.4	10.5
Cardiovascular problems	0	2.3	5.1	0	19.2*	0	36.4	15

for 16 Non Specific Health Symptoms experienced by 530 people (270 men + 260 women) in relation to their distances from cellular phone base stations (≤ 300 m vs. > 300 m [reference group]).

* = $P < 0.05$ for levels of complaints 2 + 3 pooled.

Influence of age on the percentage of complaints

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. [Investigation on the health of people living near mobile telephone relay stations: I/ Incidence according to distance and sex]. *Pathol Biol (Paris)* (2002); 50(6):369-373.

Santini R SP, Le Ruz P, Danze J, Seigne M. Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine* (2003); 22(1):41-49.

Shebeen El-Kom, Egypt (2003)



Study of 85 inhabitants living near the first cell phone tower in the city (tower operational since 1998).

37 subjects lived in the building under the transmitters.


48 others worked in the building across the street.

A control group of 80 individuals worked in a building 2 kilometers away from the towers.

Controls were matched for age, sex, occupation, education level, and mobile phone use.

Abdel-Rassoul G, El-Fateh OA, Salem MA et al. Neurobehavioral effects among inhabitants around mobile phone base stations. *Neurotoxicology* (2007); 28(2):434-440.


Shebeen El-Kom, Egypt (2003)



* = statistically significant at this sample size ($p < 0.05$ or better)

Abdel-Rassoul G, El-Fateh OA, Salem MA et al. Neurobehavioral effects among inhabitants around mobile phone base stations. *Neurotoxicology* (2007); 28(2): 434-440.

Vienna and Carinthia, Austria (2004)



Study of health effects around 10 cell phone towers (“base stations”) in urban and rural Austria.

Criteria:

Towers operational for >2 years.

No local controversy.

No other towers nearby (when possible).


900 MHz transmission.

Random selection of households within the study areas.

Performance tests, symptom questionnaires, exposure measurements in the subject’s bedroom.

Hutter HP, Moshhammer H, Wallner P, Kundi M. Subjective symptoms, sleeping problems, and cognitive performance in subjects living near mobile phone base stations. *Occup Environ Med* (2006); 63(5):307-313.

Austria (2004)



Percentage of subjects reporting symptoms, stratified by RF exposure levels as measured in subject's bedroom.

* = statistically significant for this sample size.

FCC Guidelines: 6000 mW/m²


Hutter HP, Moshammer H, Wallner P, Kundi M. Subjective symptoms, sleeping problems, and cognitive performance in subjects living near mobile phone base stations. *Occup Environ Med* (2006); 63(5):307-313.

BACKGROUND: The erection of mobile telephone base stations in inhabited areas has raised concerns about possible health effects caused by emitted microwaves.

METHODS: In a cross-sectional study of randomly selected inhabitants living in urban and rural areas for more than one year near to 10 selected base stations, 365 subjects were investigated. Several cognitive tests were performed, and wellbeing and sleep quality were assessed. Field strength of high-frequency electromagnetic fields (HF-EMF) was measured in the bedrooms of 336 households.

RESULTS: Total HF-EMF and exposure related to mobile telecommunication were far below recommended levels (max. 4.1 mW/m²). Distance from antennae was 24-600 m in the rural area and 20-250 m in the urban area. Average power density was slightly higher in the rural area (0.05 mW/m²) than in the urban area (0.02 mW/m²). Despite the influence of confounding variables, including fear of adverse effects from exposure to HF-EMF from the base station, there was a significant relation of some symptoms to measured power density; this was highest for headaches. Perceptual speed increased, while accuracy decreased insignificantly with increasing exposure levels. There was no significant effect on sleep quality. CONCLUSION: Despite very low exposure to HF-EMF, effects on wellbeing and performance cannot be ruled out, as shown by recently obtained experimental results; however, mechanisms of action at these low levels are unknown.

Akrotiri, Cyprus (2005)



Evaluation of health concerns near a military radar antenna:

Measurement of average RF levels in two nearby communities:


Akrotiri [red circle]

Asomatos [yellow circle]

And as a control, in another village > 20 km distant [blue circle]

Akrotiri also had a cell phone tower.

Akrotiri, Cyprus (2005)



On left, average RF readings in the three communities (in v/M).

On the right, percentages of four reported symptoms were significantly higher in the towns with higher RF exposures.

Average power densities:

Akrotiri: 0.57 v/m = 0.863 $\mu\text{W}/\text{cm}^2$

Asomatos: 0.46 v/m = 0.561 $\mu\text{W}/\text{cm}^2$

Pano Kyvides: 0.001 v/m = 0.000001 $\mu\text{W}/\text{cm}^2$

FCC Guidelines: 600–1000 $\mu\text{W}/\text{cm}^2$

Preece AW, Georgiou AG, Dunn EJ, Farrow SC. Health response of two communities to military antennae in Cyprus. *Occup Environ Med* (2007); 64(6):402-408.

Selbitz, Bavaria (2009)



General health survey sent to 1080 residents of the village of Selbitz, Bavaria (population 4644), with 251 responses (23% return).

Two cell tower transmitters in the center of town.

Exposure areas determined by concentric circles of 100 to 400 meters radius.


Field measurements stratified exposures into two regions:

Groups 1 and 2 (mean exposure 1.17 V/m)

Groups 3 and 4 (mean exposure 0.70 V/m)

Eger H, Jahn M. Specific Health Symptoms and Cell Phone Radiation in Selbitz (Bavaria, Germany) -- Evidence of a Dose-Response Relationship. *umwelt-medizin-gesellschaft* (2010); 23:1-20.

Selbitz, Bavaria (2009)



Some sample data from the study. Symptoms scored on 0 - 5 scale.

*** = statistically significant (Groups 1 + 2 compared to Groups 3 + 4).**

14 of 19 symptom categories showed statistically significant elevations in groups 1 and 2 as compared to groups 3 and 4.

Eger H, Jahn M. Specific Health Symptoms and Cell Phone Radiation in Selbitz (Bavaria, Germany) -- Evidence of a Dose-Response Relationship. *umwelt-medizin-gesellschaft* (2010); 23:1-20.

Danish Birth Cohort Studies



Several studies done on large numbers of young children in the Danish National Birth Cohort.

Evaluating health status of children at age 7.


Significant changes found in three areas:

Behavioral problems.

Headaches.

Hearing Loss.

Cell Phone Exposure and Behavioral Problems in Young Children



Association of prenatal and postnatal cell phone exposure and behavioral problems in young children, evaluated at age 7.

Two studies with the Danish National Birth Cohort:


Study 1 - (2006 data, n = 13,159) OR=1.80 (95% CL=1.45-2.23)

Divan HA, Kheifets L, Obel C, Olsen J. Prenatal and postnatal exposure to cell phone use and behavioral problems in children. *Epidemiology* (2008); 19(4): 523-529.

Study 2 - (2008 data, n = 28,745) OR=1.5 (95% CL=1.4-1.7)

Divan HA, Kheifets L, Obel C, Olsen J. Cell phone use and behavioural problems in young children. *J Epidemiol Community Health* (2010); 66:524-529

Cell Phone Exposure and Behavioral Problems in Young Children




Association of prenatal and postnatal cell phone exposure and behavioral problems in young children, evaluated at age 7.

Study 1 - 2006 (n = 13,159)

Divan HA, Kheifets L, Obel C, Olsen J. Prenatal and postnatal exposure to cell phone use and behavioral problems in children. *Epidemiology* (2008); 19(4): 523-529.

Behavior Problems Stratified by Cell Phone Exposure Level



Association of prenatal and postnatal cell phone exposure and behavioral problems in young children, evaluated at age 7.


Danish National Birth Cohort:

Study 2 - 2008 cohort (n = 28,745)

1 = no increased risk

Divan HA, Kheifets L, Obel C, Olsen J. Cell phone use and behavioural problems in young children. *J Epidemiol Community Health* (2010); 66:524-529

Cell Phone Exposure and Headaches in Young Children




Association of postnatal cell phone exposure and headaches in young children, evaluated at age 7.

52,680 children in the Danish National Birth Cohort:

Results—Our analyses included data from 52,680 children. Children with cell phone exposure had higher odds of migraines and headache-related symptoms than children with no exposure. The odds ratio for migraines was 1.30 (95% confidence interval: 1.01–1.68) and for headache-related symptoms was 1.32 (95% confidence interval: 1.23–1.40) for children with both prenatal and postnatal exposure.

Sudan M, Kheifets L, Arah O, Olsen J, Zeltzer L. Prenatal and Postnatal Cell Phone Exposures and Headaches in Children. *Open Pediatr Med Journal* (2012); 6(2012):46-52.

Cell Phone Use and Hearing Loss in Young Children



Association of postnatal cell phone exposure and hearing loss in young children, evaluated at age 7.

52,680 children in the Danish National Birth Cohort:

Three forms of statistical analysis showed this trend.

Results: Our analyses included data from 52 680 children. We observed weak associations between cell phone use and hearing loss at age 7, with odds ratios and 95% confidence intervals from the traditional logistic regression, MSM and DRE models being 1.21 [95% confidence interval [CI] 0.99, 1.46], 1.23 [95% CI 1.01, 1.49] and 1.22 [95% CI 1.00, 1.49], respectively.

Sudan M, Kheifets L, Arah OA, Olsen J. Cell phone exposures and hearing loss in children in the Danish National Birth Cohort. *Paediatr Perinat Epidemiol* (2013); 27(3):247-257.

Wireless “smart meters” increase public RF exposure.




In the last ten years, networks of “smart meters” have been installed in residential areas in many parts of the world.

Many of these smart meter networks operate on a “mesh” network technology where the meters transmit brief but quite strong pulses multiple times per minute, 24 hours a day.

This produces a significant increase on the total load of microwave RF in residential communities.

Onset of new symptoms after exposure to “smart meters”.




This increased residential exposure to “smart meter” RF transmissions has led to an increase in the number of individuals who report onset of symptoms of EHS.

Data from a survey conducted by Richard Conrad PhD which was presented as testimony to the Maine Public Utilities Commission.

In February 2013, the expert testimony of Richard H. Conrad, PhD, and many other experts worldwide, was submitted to the Maine Public Utilities Commission when the Commission was considering the future of Smart Meters in that state. Dr. Conrad reported the results of a survey of 210 individuals who had experienced symptoms resulting from exposure to Smart Meters. [Pre-Filed Testimony of Richard H. Conrad, Ph.D. Maine Public Utilities Commission Docket No. 2011-00262, Appendix 3 (<http://www.mainecoalitiontostopsmartmeters.org/wp-content/uploads/2013/01/Exhibit-9-Conrad-Web.pdf>).]

Use of wifi before and after “smart meter” exposure.



This chart shows the change in usage of or tolerance to wifi by respondents to Dr. Conrad’s questionnaire, before and after installation of the residential “smart meter” infrastructure.

Numbers are stated as percentages of 210 respondents.

Pre-Filed Testimony of Richard H. Conrad, Ph.D., Appendix 2 for worsened symptoms and Appendix 3 for new symptoms (<http://www.mainecoalitiontostopsmartmeters.org/wp-content/uploads/2013/01/Exhibit-9-Conrad-Web.pdf>).

Self-Reporting of Symptom Development From Exposure to Radiofrequency Fields of Wireless Smart Meters in Victoria, Australia: A Case Series

ABSTRACT

Context • In 2006, the government in the state of Victoria, Australia, mandated the rollout of smart meters in Victoria, which effectively removed a whole population's ability to avoid exposure to human-made high-frequency nonionizing radiation. This issue appears to constitute an unprecedented public health challenge for Victoria. By August 2013, 142 people had reported adverse health effects from wireless smart meters by submitting information on an Australian public Web site using its health and legal registers.

Objective • The study evaluated the information in the registers to determine the types of symptoms that Victorian residents were developing from exposure to wireless smart meters.

Design • In this case series, the registers' managers eliminated those cases that did not clearly identify the people providing information by name, surname, postal address, and/or e-mail to make sure that they were genuine registrants. Then they obtained consent from participants to have their deidentified data used to compile the data for the case series. The author later removed any individual from outside of Victoria.


Participants • The study included 92 residents of Victoria, Australia.

Outcome Measures • The author used her medical experience and judgment to group symptoms into clinically relevant clusters (eg, pain in the head was grouped with headache, tinnitus was grouped with ringing in the ears). The author stayed quite close to the wording used in the original entries. She then calculated total numbers and percentages for each symptom cluster. Percentages were rounded to the nearest whole number.

Results • The most frequently reported symptoms from exposure to smart meters were (1) insomnia, (2) headaches, (3) tinnitus, (4) fatigue, (5) cognitive disturbances, (6) dysesthesias (abnormal sensation), and (7) dizziness. The effects of these symptoms on people's lives were significant.

Conclusions • Review of some key studies, both recent and old (1971), reveals that the participants' symptoms were the same as those reported by people exposed to radiofrequency fields emitted by devices other than smart meters. Interestingly, the vast majority of Victorian cases did not state that they had been sufferers of electromagnetic hypersensitivity syndrome (EHS) prior to exposure to the wireless meters, which points to the possibility that smart meters may have unique characteristics that lower people's threshold for symptom development. (*Altern Ther Health Med.* 2014;20(6):28-39.)


Common reported symptoms from “smart meter” exposure.



This is a list of the most common symptoms reported as provoked from “smart meter” exposure in the Victoria, Australia study, graphed as percent of 92 respondents who reported the symptom.

Lamech, F. Self-Reporting of Symptom Development From Exposure to Radiofrequency Fields of Wireless Smart Meters in Victoria, Australia: A Case Series. *Alternative Therapies*, 2014 (Nov/Dec) Vol. 20:6.


Common reported symptoms from “smart meter” exposure.



This is a list of the most common symptoms reported as provoked from “smart meter” exposure in the Victoria, Australia study, graphed as percent of 92 respondents who reported the symptom.

Lamech, F. Self-Reporting of Symptom Development From Exposure to Radiofrequency Fields of Wireless Smart Meters in Victoria, Australia: A Case Series. *Alternative Therapies*, 2014 (Nov/Dec) Vol. 20:6.


Common reported symptoms from “smart meter” exposure.



This is a list of the most common symptoms reported as provoked from “smart meter” exposure in the Victoria, Australia study, graphed as percent of 92 respondents who reported the symptom.

Lamech, F. Self-Reporting of Symptom Development From Exposure to Radiofrequency Fields of Wireless Smart Meters in Victoria, Australia: A Case Series. *Alternative Therapies*, 2014 (Nov/Dec) Vol. 20:6.


Wifi networks also produce constant exposure to RF transmissions.



Wifi routers transmit constantly, whether or not they are being “used” by the computers in the network.

We are now in the process of putting this sort of constant RF exposure into all our schools, and all of our public spaces.

Exposure to microwave RF has increased dramatically.




Measurement of average levels of microwave RF signal (900 MHz - 2.5 GHz) in urban areas in France, 2000-2012.

The green line is at 0.1 microwatt/cm² = 1000 microwatts per meter squared ~ 0.6 V/m, which many researchers in Europe suggest as an appropriate limit for exposure to the public.

<http://www.next-up.org>

Incidence of self-identified electrohypersensitivity (%)



In the previous studies, we saw that some symptoms are more common with higher exposure to microwave RF transmissions.


The people who had those symptoms may or may not have been aware that RF was a factor.

But some people with symptoms triggered by microwave RF exposures are aware that this is happening to them.

If these symptoms are sufficiently debilitating, the affected individual may consider themselves to be “electrohypersensitive”.

The above graph shows the percentage of the population that self-identified as having “EHS” in surveys done in various countries over the last two decades.

The prevalence of self-reported EHS is increasing.



The prevalence (%) of people around the world who consider themselves to be electrosensitive, as reported in various research studies from 1997 to 2008, plotted over time in a normal distribution graph.

The endpoint at 50% is an extrapolated value.

Hallberg O, Oberfeld G. Letter to the editor: will we all become electrosensitive? *Electromagn Biol Med* (2006); 25(3): 189-191.

Stockholm County, Sweden, 1997: 1.5% of the population reported being hypersensitive to electrical or magnetic fields. (Hillert et al., 2002)

California, 1998: 3.2% of the adult population reported being sensitive to sources of EMF. (Levallois et al., 2002)

Switzerland, 2004: 5% of the population had symptoms attributable to EHS. (Schreier et al., 2006)

Austria, 2004: 2% of the population was estimated to have electrohypersensitivity.

Austria, 2008: 29.3% with some adverse response, 2.1% reported intense disturbance, and 3.5% had experienced enough difficulty that they had consulted a physician about the problem. (Schrottner and Leitgeb, 2008)

Europe is ahead of us on the RF learning curve.



The evidence on acute effects and on long-term degenerative changes produced by chronic RF exposure will continue to build over the next few years.

Public concern is likely to continue to grow over time, especially as regards the exposure of children in an environment where their attendance is required by law.

These problems and concerns will have legal repercussions, down the line.

One way to get perspective on this problem is to observe what is happening in Europe, where the wireless revolution started, and the public has been exposed for a longer interval of time.

2008 – French National Library removes WiFi.

La Bibliothèque Nationale de France renonce au WiFi.

Paris 07 04 2008 - La direction de la célèbre Bibliothèque Nationale de France (BNF) vient de décider un moratoire sur les Hots Spots WiFi d'accès à l'internet qui devaient être installés par une société privée sur l'ensemble de la zone.

Les motifs évoqués étant la recherche de la qualité du service, mais aussi l'application du principe de précaution afin de ne pas exposer son personnel et l'ensemble des visiteurs aux risques des irradiations par les champs électromagnétiques de 2,4 GHz.

Cette décision est fondée sur une argumentation qui s'appuie sur la littérature scientifique qui prouve les effets génotoxiques des ondes du WiFi, notamment :
- l'étude scientifique américaine de l'université de Chicago de l'équipe du Pr S. Lee concernant les modifications génétiques des cellules humaines exposées aux champs des RadioFréquences de 2,45 GHz (fréquence du WiFi).

[Etude scientifique : [NCBI.Gov – PubMed](#)]

- Le rapport du [Consortium Scientifique BioInitiative](#) qui conclut à la révision immédiate des normes actuelles de l'exposition des populations aux ondes électromagnétiques, [[Pétition Internationale](#) en cours], ainsi que [l'étude scientifique](#) de l'université Blaise Pascal de Clermont-Ferrand.

Arnaud Beaufort, directeur général adjoint de la BNF confirme le moratoire en ces termes :

" ... nous attendons les résultats d'études en cours et nous nous donnons le temps de choisir la technologie la plus adaptée."



La Bibliothèque Nationale de France [en chiffres](#) :
- la BNF comprend dans les 4 tours développant plus de 200 000 m² de surface un fond de quinze millions d'ouvrages divers en livres, manuscrits, estampes, dessins, photographies, etc . . . pour une fréquentation de plus d'un million de personnes/an avec un personnel comprenant plus de 2500 collaborateurs et un budget de 254 M€/an.

The Bibliotheque Nationale de France eliminated WiFi from the library in 2008.

This came about in part because growing numbers of library staff were developing symptoms of EHS.

In 2009, three of the major university libraries in Paris followed suit.

France – Growing Concerns with WiFi in the Workplace.

"Après les Bibliothèques et la RATP, c'est au tour des policiers « d'en avoir par-dessus la tête » du Wifi et des antennes-relais !" - L'indépendant de Perpignan - 04/10/2012

De plus en plus de syndicats saisissent les CHS pour connaître l'impact des ondes sur la santé



Accueil



Envoyer à un ami



Version imprimable



Partager

Souvenons-nous : En 2007, un certain nombre de bibliothécaires parisiens se plaignaient de maux de têtes, nausées... après l'installation massive de bornes Wifi dans leurs établissements. Cet épisode avait abouti à un bras de fer entre la Mairie de Paris et les syndicats (donnant lieu notamment à l'organisation d'une Conférence de Citoyens sur la prolifération des ondes électromagnétiques à Paris).

A la BnF, et plus récemment à la RATP, c'est dans les mêmes (mauvaises) conditions que le débat a eu lieu entre syndicats et employeurs sur ces questions de santé au travail, conséquence du déferlement des nouvelles technologies sans fil dans l'entreprise.

In France, labor unions are more influential than they are in the United States.

Increasing concern is being expressed by labor unions regarding the health effects of WiFi in the workplace.

This started in the libraries in Paris, in 2007.

Then the labor unions in the municipal transit systems raised the issue.

By 2012, the unions of police and customs agencies were raising the same issues regarding microwave RF exposure from WiFi and cellular transmission towers located on the jobsite.

La loi sur les ondes électromagnétiques en 5 questions

Le Monde.fr | 24.01.2014 à 16h14 • Mis à jour le 24.01.2014 à 18h46 |

Par [Audrey Garric \(/journaliste/audrey-garric/\)](#) et [Pierre Le Hir \(/journaliste/pierre-le-hir/\)](#)



On January 23, 2013, the French parliament passed a new law covering exposure to microwave RF.

This law prohibits the operation of WiFi in childcare settings for children of age 3 years or less.

It prohibits the telcom industry from directing any advertising to the demographic ≤ 14 years of age.

It requires vendors of cell phones to provide a hands-free option for the phone.

The French administration is also planning to make a proposal to parliament to address the possibility of creating zones of limited EMF exposure within French cities, to address the problems of individuals with EHS.

Hamburg, Germany has banned wifi from schools.

Hamburg legt Pläne für WLAN an Schulen auf Eis

vom 29. November 2014

Aus der Redaktion des Flensburger Tageblatts

An sechs Hamburger Schulen sollte das digitale Zeitalter Einzug halten. Stadt und Schulbehörde rudern zurück: Nun werde geprüft, ob das Funknetz WLAN zu gefährlich für die Gesundheit der Schüler ist.

Empfehlen 289

Twittern 20

Pin it

g+1 379

Kommentare 32



Lernen mit dem Tablet: Das wird es in Hamburg so bald nicht geben. Zu groß sind die Bedenken bezüglich der Gesundheit der Schüler.

At six schools in Hamburg the digital age should find their way. City and school board now back row: The wireless network WLAN was too dangerous for the health of students.

<http://www.shz.de/nachrichten/deutschland-welt/netzwelt/hamburg-legt-plaene-fuer-wlan-an-schulen-auf-eis-id8321086.html>

Israeli Supreme Court – 2013

Israeli Supreme Court Ordered Israeli Government to Investigate Number of Children Suffering From EHS

Dafna Tachover, Jul 22, 2013

On July 18, 2013, the Israeli Supreme Court ordered the Israeli Government to investigate how many children in Israel already suffer from electromagnetic hypersensitivity (EHS). The Government must submit the result of its investigation to the court by November 16, 2013.

The Order was issued in response to a Petition to the Israeli Supreme Court to ban Wi-Fi in schools submitted in August, 2012.

This order is the result of an extensive brief written by Attorney Dafna Tachover and submitted to the Court on June 13, 2013. A 65-page brief, supported by 640 pages of appendices, emphasized the problem of EHS and claimed that it is unreasonable to expose children to WiFi when it is proven to cause sickness.

The EU is beginning to address EHS as a public health problem.

PRELIMINARY DRAFT OPINION

of the

Section for Transport, Energy, Infrastructure and the Information Society

on

Electromagnetic hypersensitivity

(own-initiative opinion)

Rapporteur: **Bernardo Hernández Bataller**

To the members of the Study Group on **Electromagnetic hypersensitivity**
(Section for Transport, Energy, Infrastructure and the Information Society)


N.B.: This document will be discussed at the meeting on **5 December 2014** beginning at **9.30 a.m.**

European Economic and Social Committee – TEN/559 Electromagnetic Hypersensitivity

“Electromagnetic hypersensitivity syndrome is a complex problem which has to be solved through a combination of legislative and other measures. In terms of fundamental rights, there is a conflict between the rights of sufferers, their physical integrity and health, on the one hand, and the right to freedom of communication, on the other hand. These rights need to be taken into account before legislation can be adopted on this issue. The EESC is in favour of adopting binding legislation that reduces or mitigates exposure to electromagnetic fields.

“The EU should assist currently affected groups and limit exposure fields in light of the recommendations set out in this opinion, especially with respect to recognising this exposure as a cause of functional disability. Steps should also be taken to prevent the number of sufferers from gradually increasing in the future due to the expansion of devices using these technologies.”

Alterations in Hormone Physiology



Now we turn to a discussion of the research showing effects of microwave RF on hormone physiology.

Rimbach, Bavaria (2004 - 2005)



In spring of 2004 a GSM cell tower was installed near Rimbach, Bavaria (population ~ 2000).


Prior to activation of the antenna, the town residents were asked to participate in a mass screening.

Urine levels of the stress hormones adrenaline, noradrenaline, dopamine, and phenylethylamine were measured in January/February 2004, and again in July 2004, January 2005, and July 2005.

A medical history and symptom questionnaire was also administered.

Buchner K EH. Changes of Clinically Important Neurotransmitters under the Influence of Modulated RF Fields--A Long-term Study under Real-life Conditions. *Umwelt-Medizin-Gesellschaft* (2011); 24(1):44-57.

Rimbach, Bavaria (2004 - 2005)




Here we see some symptom scores before tower activation (blue) and after a year of tower transmission (red)
Some health complaints increased over the course of the study.

Buchner K EH. Changes of Clinically Important Neurotransmitters under the Influence of Modulated RF Fields--A Long-term Study under Real-life Conditions. *Umwelt-Medizin-Gesellschaft* (2011); 24(1):44-57.

Abstract: This follow-up of 60 participants over one and a half years shows a significant effect on the adrenergic system after the installation of a new cell phone base station in the village of Rimbach (Bavaria). After the activation of the GSM base station, the levels of the stress hormones adrenaline and noradrenaline increased significantly during the first six months; the levels of the precursor dopamine decreased substantially. The initial levels were not restored even after one and a half years. As an indicator of the dysregulated chronic imbalance of the stress system, the phenylethylamine (PEA) levels dropped significantly until the end of the study period. The effects showed a dose-response relationship and occurred well below current limits for technical RF radiation exposures. Chronic dysregulation of the catecholamine system has great relevance for health and is well known to damage human health in the long run.

Adrenaline levels



Results were stratified by in-home exposure levels ($\mu\text{W}/\text{m}^2$) in three cohorts.


Hormone levels graphed for each exposure cohort.

Levels of the stress hormone adrenaline rose after the transmitter became active.

In the highest exposure cohort adrenaline levels never returned to pre-exposure baseline.

Fig. 3: Median adrenaline levels for all participating citizens of Rimbach whose cell phone base station exposure was above $100 \mu\text{W}/\text{m}^2$, between 60 and $100 \mu\text{W}/\text{m}^2$, or up to $60 \mu\text{W}/\text{m}^2$. The power density levels refer to peak values of the GSM radiation exposure in front of a given residence.

Noradrenaline levels



Noradrenaline levels also rose after the transmitter became active.

They never returned to pre-exposure baseline.

Fig. 7: Median noradrenaline levels in all participating citizens of Rimbach as a function of GSM power density levels (peak values)

Effect of cordless DECT phones.





Fig. 8: Median noradrenaline values for subjects who had a DECT phone or other wireless devices at home, for those without indoor wireless devices, as well as for subjects without indoor wireless devices and with a GSM radiation exposure up to $60 \mu\text{W}/\text{m}^2$ (peak value measured in front of residence)

Buchner K EH. Changes of Clinically Important Neurotransmitters under the Influence of Modulated RF Fields--A Long-term Study under Real-life Conditions. *Umwelt-Medizin-Gesellschaft* (2011); 24(1):44-57.

Abstract: This follow-up of 60 participants over one and a half years shows a significant effect on the adrenergic system after the installation of a new cell phone base station in the village of Rimbach (Bavaria). After the activation of the GSM base station, the levels of the stress hormones adrenaline and noradrenaline increased significantly during the first six months; the levels of the precursor dopamine decreased substantially. The initial levels were not restored even after one and a half years. As an indicator of the dysregulated chronic imbalance of the stress system, the phenylethylamine (PEA) levels dropped significantly until the end of the study period. The effects showed a dose-response relationship and occurred well below current limits for technical RF radiation exposures. Chronic dysregulation of the catecholamine system has great relevance for health and is well known to damage human health in the long run.

Melatonin




The pineal gland secretes melatonin. Ambient light suppresses melatonin secretion. So melatonin secretion is high during the night-time hours, peaking shortly after midnight. **Higher melatonin levels are part of what makes us feel “sleepy” at night.**

Exposure to light during the night-time hours will lead to a rapid suppression of melatonin secretion by the pineal gland, and this can cause disruption of sleep and derangement of the circadian rhythm.

Melatonin is one of the most potent anti-oxidant molecules in the human body, and acts to reduce reactive oxidative processes in the body. Melatonin can quench the damaging free radical activity produced by inflammation. The presence of elevated melatonin at night is therefore a key factor in the healing and rejuvenating functions that we associate with “a good night’s sleep”.

Melatonin lowers risk of breast cancer.



[Figure 1. Smoothing spline plot for aMT6s level (ng/mg creatinine) in relation to breast cancer risk among postmenopausal women. 95% CIs are indicated by dotted lines.]


Melatonin is also protective against the growth of cancer cells, and disruption of the circadian melatonin cycle has been shown to lead to increased tumor growth in a variety of cancer types.

Women who have lower levels of nocturnal melatonin are at greater risk for developing breast cancer.

In 2007 the International Agency for Research on Cancer declared night shift work to be a probable carcinogen due to increased breast cancer risk..

Schernhammer ES, Hankinson SE. Urinary melatonin levels and postmenopausal breast cancer risk in the Nurses' Health Study cohort. *Cancer Epidemiol Biomarkers Prev* (2009); 18(1):74-79.

Schwarzenburg Short Wave Radio Broadcast Tower – 1998



RF exposure can also lower melatonin levels.


Schwarzenburg experiment: Decommissioning the Swiss national short-wave radio transmitter of Schwarzenburg, about 20 km south of the Swiss Capital city of Berne, transmitting since 1939. It operated at frequencies of 3 to 30 MHz, with maximum power of two times 150 kW.

Figure 1. Map of the Schwarzenburg area showing the location of the transmitter, the H-field measurement points and the location of the zones A, B, C and R. The diameters of the circles around the measurement points indicate the 24 hour average magnetic field strengths, as measured between August 1992 and August 1993. (Reproduced with approval from swisstopo (BA046633).)

Abelin T, Altpeter E, Roosli M. Sleep Disturbances in the Vicinity of the Short-Wave Broadcast Transmitter Schwarzenburg. *Somnologie* (2005); 9:203-209.

Altpeter ES, Roosli M, Battaglia M, Pfluger D, Minder CE, Abelin T. Effect of short-wave (6-22 MHz) magnetic fields on sleep quality and melatonin cycle in humans: Schwarzenburg shut-down study. *Bioelectromagnetics* (2006); 27(2):142-150.


Melatonin Excretion (pg/ml)



Altpeter ES, Roosli M, Battaglia M, Pflugger D, Minder CE, Abelin T. Effect of short-wave (6-22 MHz) magnetic fields on sleep quality and melatonin cycle in humans: the Schwarzenburg shut-down study. *Bioelectromagnetics* (2006); 27(2):142-150.


Morning Fatigue (0 - 100 Scale)

■ Baseline ■ After Shutdown



Altpeter ES, Roosli M, Battaglia M, Pfluger D, Minder CE, Abelin T. Effect of short-wave (6-22 MHz) magnetic fields on sleep quality and melatonin cycle in humans: the Schwarzenburg shut-down study. *Bioelectromagnetics* (2006); 27(2):142-150.

Microwave RF Interacts with Molecular Structures



(Original art by Raychel Ciemma, Springfield, Oregon, for Paul Lee DO FAAO's book [Interface](#), used with permission.)


The molecules in our bodies vary in size and in total electric charge.

These molecular structures of our body can resonate with fluctuating electromagnetic fields.

Any charged particle has a resonant frequency.

This frequency varies depending on the total mass and charge of the particle.

Molecules resonate in fluctuating electromagnetic fields.



The molecules in our bodies vary in size and total electric charge.


These molecular structures of our body resonate with fluctuating electromagnetic fields.

Resonance Frequency



When you push something at its resonant frequency, a small force can produce a lot of motion.

Resonance Frequency




Placing the particle in an electromagnetic field that fluctuates at the resonant frequency will amplify the motion of the particle.

This is how a cyclotron works, and the frequency is often referred to as the “Ion Cyclotron Resonance” or ICR frequency.

Magnetic fields that fluctuate at the resonant frequency of an ion like calcium, or of a specific enzyme, can have dramatic effects on biochemical processes in the body.


Resonance Effect



Planaria exposed to a magnetic field fluctuating at the calcium ion's ICR frequency take far longer (48 hours) to regenerate than those that are not exposed.

Liboff A. Weak low-frequency electromagnetic fields are biologically interactive. In: Giuliani L, Soffritti M, eds. Non-Thermal Effects and Mechanisms of Interaction Between Electromagnetic Fields and Living Matter -- An ICEMS Monograph. Fidenza, Italy: Mattioli, (2010): 51-61. <http://www.ramazzini.it/ricerca/publications.asp>

“frequency window”




Some effects of fluctuating EMF occur at specific frequencies, called “frequency windows”.

The peak in IGF-II expression for human osteosarcoma bone cells exposed to combined magnetic fields occurs when the field is tuned to the calcium ion’s ICR frequency

Liboff A. Weak low-frequency electromagnetic fields are biologically interactive. In: Giuliani L, Soffritti M, eds. Non-Thermal Effects and Mechanisms of Interaction Between Electromagnetic Fields and Living Matter -- An ICEMS Monograph. Fidenza, Italy: Mattioli, (2010): 51-61. <http://www.ramazzini.it/ricerca/publications.asp>

“power window”



At a given frequency, some power levels may have a different effect than others. This is a “power window”


In this illustration, the odds ratio for childhood onset of Acute Lymphoblastic Leukemia is significantly higher if they are exposed to 60 cycle magnetic fields at a magnitude of 0.4 to 0.499 microtesla.

Lower and higher field magnitudes do not show the same effect.

Liboff A. Weak low-frequency electromagnetic fields are biologically interactive. In: Giuliani L, Soffritti M, eds. Non-Thermal Effects and Mechanisms of Interaction Between Electromagnetic Fields and Living Matter -- An ICEMS Monograph. Fidenza, Italy: Mattioli, (2010): 51-61. <http://www.ramazzini.it/ricerca/publications.asp>

Fig. 1. Odds ratios for childhood ALL, determined by Linet et al 6, as a function of residential magnetic field. The large ratios seen for fields between .4 and .499 μT , although having many less participants, are nevertheless statistically significant.

“power window”




Mice with Ascites Ehrlich carcinoma 33,
exposed to a fluctuating EM field tuned to the ICR frequency for aspartic acid and glutamic acid ions.
Survival varies with the AMPLITUDE (magnitude) of the field.

Liboff A. Weak low-frequency electromagnetic fields are biologically interactive. In: Giuliani L, Soffritti M, eds. Non-Thermal Effects and Mechanisms of Interaction Between Electromagnetic Fields and Living Matter -- An ICEMS Monograph. Fidenza, Italy: Mattioli, (2010): 51-61. <http://www.ramazzini.it/ricerca/publications.asp>

Fig. 5. Survival curve for mice infected with Ascites Ehrlich carcinoma33, under ICR conditions corresponding to mean tuning (4.4 Hz) for aspartic acid and glutamic acid ions. In contrast to Fig. 2 where the frequency is varied, a resonance (or window) peak is observed as the AC magnetic field intensity is varied

What does this mean?



(Original art by Raychel Ciemma, Springfield, Oregon, for Paul Lee DO FAAO's book [Interface](#), used with permission.)


There are thousands of enzymes and other molecules in the human body.

Each has its own mass, charge, and resonant frequency.


This means that different electromagnetic frequencies will resonate with different molecules.

Which means that the biological effects of EMF on molecular physiology are probably much more complex than is generally assumed to be the case.

Microwave RF
Produces
Oxidative
Stress in Cells



Electromagnetic Spectrum



Ionizing radiation from the high energy end of the electromagnetic spectrum can directly break DNA molecular bonds, causing mutations.

But photons of microwave RF do not have enough energy to directly break covalent molecular bonds.

Industry advocates often make the statement that since RF cannot break molecular bonds, there is no way that it can cause cancer.

Such statements sound like good physics. But they reflect a poor understanding of biology.

Tobacco can cause cancer. Genital warts can cause cancer. Asbestos can cause cancer. There are many ways to cause cancer besides ionizing radiation.

Free Radicals

**A normal molecule has
an even number of
“paired” electrons**



**A free radical molecule
has an odd number -
An “unpaired” electron**



Free radicals are oxidizing agents.

They take electrons from other atoms or molecules, which can break molecular bonds.

Life cycle of a free radical.

initiation



propagation




termination



Precursor molecule (AB) splits to form two free radicals.

Free radicals then can produce chain reactions, causing oxidative damage.



Peroxisomes (yellow) in a cell - packages of free radicals stored in cells.

Cells are making free radicals all the time.

Our bodies release them in inflammation to combat bacteria, remove diseased tissue, etc.

The free radicals release by the inflammatory process can break covalent bonds and fragment macromolecules.

Guest Editor:

Electromagnetic fields act *via* activation of voltage-gated calcium channels to produce beneficial or adverse effects

Martin L. Pall *

Professor Emeritus of Biochemistry and Basic Medical Sciences, Washington State University, Portland, OR, USA

Received: January 8, 2013; Accepted: May 20, 2013

- Introduction
- Possible modes of action following voltage-gated calcium channel stimulation
- Therapeutic bone-growth stimulation *via* Ca²⁺/nitric oxide/cGMP/protein kinase G
- Ca²⁺/nitric oxide/peroxynitrite and pathophysiological responses to EMF exposures: the example of single-strand DNA breaks
- Discussion and conclusions

Abstract

The direct targets of extremely low and microwave frequency range electromagnetic fields (EMFs) in producing non-thermal effects have not been clearly established. However, studies in the literature, reviewed here, provide substantial support for such direct targets. Twenty-three studies have shown that voltage-gated calcium channels (VGCCs) produce these and other EMF effects, such that the L-type or other VGCC blockers block or greatly lower diverse EMF effects. Furthermore, the voltage-gated properties of these channels may provide biophysically

This recently published article reviews published evidence that EMF can produce physiologic effects by altering the function of voltage gated calcium channels in cell walls.

Pall ML. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. *J Cell Mol Med* (2013);

Table 1 EMF responses blocked or lowered by calcium channel blockers

Ref. no.	EMF type	Calcium channel	Cell type or organism	Response measured
2	Pulsed magnetic fields	L-type	Human lymphocytes	Cell proliferation; cytokine production
3	Static magnetic field (0.1 T)	L-type	Human polymorphonuclear leucocytes	Cell migration; degranulation
5	ELF	L-type	Rat chromaffin cells	Differentiation; catecholamine release
6	Electric field	L-type	Rat and mouse bone cells	Increased Ca ²⁺ , phospholipase A2, PGE2
7	50 Hz	L-type	Mytilus (mussel) immunocytes	Reduced shape change, cytotoxicity
8	50 Hz	L-type	AtT20 D16V, mouse pituitary corticotrope-derived	Ca ²⁺ increase; cell morphology, premature differentiation
9	50 Hz	L-type	Neural stem/progenitor cells	<i>In vitro</i> differentiation, neurogenesis
10	Static magnetic field	L-type	Rat	Reduction in oedema formation
11	NMR	L-type	Tumour cells	Synergistic effect of EMF on anti-tumour drug toxicity
12	Static magnetic field	L-type	Myelomonocytic U937 cells	Ca ²⁺ influx into cells and anti-apoptotic effects
13	60 Hz	L-type	Mouse	Hyperalgesic response to exposure
14	Single nanosecond electric pulse	L-type	Bovine chromaffin cells	Very rapid increase in intracellular Ca ²⁺

These are some of the 23 published studies documenting that EMF can increase flow through these calcium channels, producing biological effects.

In all these studies, the effects of EMF on increased cellular calcium levels could be blocked by calcium channel blocking drugs.

Pall ML. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. *J Cell Mol Med* (2013);


15	Biphasic electric current	L-type	Human mesenchymal stromal cells	Osteoblast differentiation and cytokine production
16	DC & AC magnetic fields	L-type	β -cells of pancreas, patch clamped	Ca ²⁺ flux into cells
17	50 Hz	L-type	Rat pituitary cells	Ca ²⁺ flux into cells
18	50 Hz	L-type, N-type	Human neuroblastoma IMR32 and rat pituitary GH3 cells	Anti-apoptotic activity
19	Nanosecond pulse	L-type, N-type, P/Q-type	Bovine chromaffin cells	Ca ²⁺ dynamics of cells
20	50 Hz	Not determined	Rat dorsal root ganglion cells	Firing frequency of cells
21	700–1100 MHz	N-type	Stem cell–derived neuronal cells	Ca ²⁺ dynamics of cells
22	Very weak electrical fields	T-type	Sharks	Detection of very weak magnetic fields in the ocean
23	Short electric pulses	L-type	Human eye	Effect on electro-oculogram
24	Weak static magnetic field	L-type	Rabbit	Baroreflex sensitivity
25	Weak electric fields	T-type	Neutrophils	Electrical and ion dynamics
26	Static electric fields, 'capacitive'	L-type	Bovine articular chondrocytes	Agrican & type II collagen expression; calcineurin and other Ca ²⁺ /calmodulin responses

EMF: electromagnetic field; ELF: extremely low frequency.

In all these studies, the effects of EMF on increased cellular calcium levels could be blocked by calcium channel blocking drugs.

Pall ML. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. *J Cell Mol Med* (2013);

EMF Activation of VGCCs Increases Free Radical Production



Normally, Calcium concentrations are much higher outside of cells than inside them.

Influxes of calcium into cells act as chemical signals to alter cellular physiologic activity.


Here we have a diagram of a cell, with high levels of calcium outside, and lower levels of calcium inside.

The green arrow is a voltage-gated calcium channel, that can open to allow more calcium to enter the cell.

Inside the cell, we can see an enzyme (nitric oxide synthetase).


As discussed by Pall ML. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. *J Cell Mol Med* (2013);

EMF Activation of VGCCs Increases Free Radical Production



An electromagnetic field arrives at the cell wall.


EMF Activation of VGCCs Increases Free Radical Production



The electromagnetic field stimulates opening of voltage-gated calcium channels (VGCCs) in the cell membrane.


This increases Ca^{++} entry into the cell.

EMF Activation of VGCCs Increases Free Radical Production




Increased intracellular calcium levels stimulate the activity of nitric oxide synthetase,
Which leads to increased production of nitric oxide in the cell.

EMF Activation of VGCCs Increases Free Radical Production



Increased nitric oxide leads to increase in peroxynitrite, a potent non-radical oxidant.

EMF Activation of VGCCs Increases Free Radical Production




Peroxynitrite produces free radicals, including hydroxyl radical and NO₂.

This increase in free radicals then leads to inflammation, oxidant stress, and damage to cell structures, including DNA.

The EMF doesn't directly damage the cell. It just deranges cellular metabolism.

The free radicals that are produced by this change in metabolism are what causes the damage.

Oxidative Stress From Microwave RF Damages DNA




The mechanisms of **how** RF increases free radical activity and oxidative stress are still being explored.

But the fact that RF **does do this** has been CLEARLY ESTABLISHED by many research studies.

This increase in free radical levels can and does lead to DNA damage.

Comet assay: Unexposed control



The Comet assay is one way to measure DNA damage.

This is a study of DNA extracted from normal rat brain cells (unexposed controls).

Electrophoresis: DNA molecules of given mass and charge placed in a diffusion medium.

Preparation placed in a static electric field.

DNA molecules migrate towards a charged pole.

DNA molecules that are the same size, so they migrate at the same rate, will stay in a clump.

Kesari KK, Behari J, Kumar S. Mutagenic response of 2.45 GHz radiation exposure on rat brain. *Int J Radiat Biol* (2010a); 86(4):334-343.


RF exposure: 2.45 GHz @ 0.34 mW/cm², 2 hours per day x 35 days



DNA from brain cells of exposed rats. Here, some of the DNA molecules are broken.
The broken parts vary in mass and total charge, so they migrate through the gel at different rates
This leaves a “comet tail” of lighter fragments behind the main body of intact DNA.
The length of the tail can be measured.
This is a **very sensitive** assay for DNA damage.

Kesari KK, Behari J, Kumar S. Mutagenic response of 2.45 GHz radiation exposure on rat brain. *Int J Radiat Biol* (2010a); 86(4):334-343.

RF exposure: 2.45 GHz @ 0.34 mW/cm², 2 hours per day x 35 days




Comet Assay: **Measure of DNA fragmentation** in rat brains, produced by prolonged exposure to microwave RF.

In this study, exposure was 2 h a day for 35 days an exposure level of **one third of the FCC exposure limit**.

FCC exposure limit = 1 mW/cm²

Kesari KK, Behari J, Kumar S. Mutagenic response of 2.45 GHz radiation exposure on rat brain. *Int J Radiat Biol* (2010a); 86(4):334-343.

RF exposure: 2.45 GHz @ 0.34 mW/cm², 2 hours per day x 35 days



Depletion of antioxidants in RF-exposed rat brains.


This consumption of anti-oxidants is **evidence of increased oxidant stress**, due to excess free radical production.

Kesari KK, Behari J, Kumar S. Mutagenic response of 2.45 GHz radiation exposure on rat brain. *Int J Radiat Biol* (2010a); 86(4):334-343.

Abstract

Purpose: To investigate the effect of 2.45 GHz microwave radiation on rat brain of male wistar strain. **Material and methods:** Male rats of wistar strain (35 days old with 130 + 10 g body weight) were selected for this study. Animals were divided into two groups: Sham exposed and experimental. Animals were exposed for 2 h a day for 35 days to 2.45 GHz frequency at 0.34 mW/cm² power density. The whole body specific absorption rate (SAR) was estimated to be 0.11 W/Kg. Exposure took place in a ventilated Plexiglas cage and kept in anechoic chamber in a far field configuration from the horn antenna. After the completion of exposure period, rats were sacrificed and the whole brain tissue was dissected and used for study of double strand DNA (Deoxyribonucleic acid) breaks by micro gel electrophoresis and the statistical analysis was carried out using comet assay (IV-2 version software). Thereafter, antioxidant enzymes and histone kinase estimation was also performed. **Results:** A significant increase was observed in comet head (P50.002), tail length (P50.0002) and in tail movement (P50.0001) in exposed brain cells. An analysis of antioxidant enzymes glutathione peroxidase (P50.005), and superoxide dismutase (P50.006) showed a decrease while an increase in catalase (P50.006) was observed. A significant decrease (P50.02) in histone kinase was also recorded in the exposed group as compared to the control (sham-exposed) ones. One-way analysis of variance (ANOVA) method was adopted for statistical analysis. **Conclusion:** The study concludes that the chronic exposure to these radiations may cause significant damage to brain, which may be an indication of possible tumour promotion (Behari and Paulraj 2007).

RF exposure: 2.45 GHz @ 0.21 mW/cm², 2 hours per day x 45 days




Suppression of melatonin secretion by 2.45 GHz RF:

Bad news, since melatonin is also a potent antioxidant.

Kesari KK, Kumar S, Behari J. Pathophysiology of microwave radiation: effect on rat brain. *Appl Biochem Biotechnol* (2012); 166(2):379-388.

Evidence of DNA damage by microwave RF.




Another study, using Human fibroblasts.
1950 MHz, 5 minutes on/10 minutes off.
Total exposure for 4, 8, or 24 hours.
DNA fragmentation measured by Comet Assay.

Figure 9. Intermittent RF-EMF exposure (1950 MHz, 5 minutes on/10 minutes off, 1 and 2 W/kg, 4, 8 and 24 hours) increases the DNA strand break frequency in human fibroblasts dependent on the duration of exposure as measured with the alkaline and neutral Comet assay (H.-W. Rüdiger et al., Division of Occupational Medicine, University of Vienna, Austria).

Adlkofer F. Risk Evaluation of Potential Environmental Hazards from Low Energy Electromagnetic Field Exposure Using Sensitive In Vitro Methods. *Bioelectromagnetics* (2006); 331-354.

DNA damage blocked by anti-oxidants



A cell study, with human fibroblasts, exposed to 1950 MHz RF, 5 minutes on/10 minutes off.

(right hand columns => **DNA damage blocked by anti-oxidant** effect of vitamin C (ascorbic acid)).

The research group of Prof. Tauber, Berlin, investigated the effect of RF-EMF on HL-60 cells, i.e. a human promyelocytic cell line. After continuous exposure to RF-EMF of 1800 MHz and a SAR value of 1.3 W/kg they observed a highly significant increase in the number of single and double DNA strand breaks as measured by the alkaline Comet assay and of micronuclei as measured with the micronucleus test, thus fully confirming the findings obtained in the Vienna laboratory. Additionally, as clearly shown in Figures 12 and 13, the generation of DNA strand breaks and micronuclei can be prevented, when the radical scavenger ascorbic acid is added to the culture medium before exposure.

Figure 12, from: Adlkofer F. Risk Evaluation of Potential Environmental Hazards from Low Energy Electromagnetic Field Exposure Using Sensitive In Vitro Methods. *Bioelectromagnetics* (2006); 331-354.


Consequences of Chronic Microwave RF Exposure



Fig. 2. Geographical location of BS Site BH 20 at 1373 Rua do Ouro Street, in the Serra neighborhood, Belo Horizonte municipality

Dode AC, Leao MM, Tejo Fde A et al. Mortality by neoplasia and cellular telephone base stations in the Belo Horizonte municipality, Minas Gerais state, Brazil. *Sci Total Environ* (2011); 409(19):3649-3665.

Amateur Radio Operators




Analysis of leukemia deaths in male members of the American Radio Relay League resident in Washington and California, 1971-1983

Milham SJ. Silent keys: leukaemia mortality in amateur radio operators. *Lancet* (1985); 1(8432):812.

Cherry N. Evidence in support of the a priori hypothesis that Electromagnetic Radiation across the spectrum is a Ubiquitous Universal Genotoxic Carcinogen. (2002):1-52. <http://www.neilcherry.com/documents.php>

Polish Military (1971-1985)



Polish military personnel with occupational exposure to radio and microwave frequency radiation.

Odds ratio of cancer incidence (1971-1985)


CML = chronic myelocytic leukemia

AML = acute myeloblastic leukemia

NHL = non-Hodgkin lymphoma

Szmigielski S. Cancer morbidity in subjects occupationally exposed to high frequency (radiofrequency and microwave) electromagnetic radiation. *Sci Total Environ* (1996); 180(1):9-17.

U.S. Navy Korean War Veterans (1950-1974)



Mortality in U.S. Navy Korean War Veterans (1950–1974) stratified by levels of occupational radar exposure.

Mortality 1950–1974. (Y axis = crude mortality per 1000) Stratified by level of radar exposure.


In the original paper, Robinette et al evaluated job exposure hazard levels of 6 categories of navy personnel and grouped them into two groups, low exposure and high exposure. The electronic technicians (ET) had a significantly lower hazard rating and lower levels of pathology than the other two job categories in the high risk group, so this classification diluted out the high exposure risk pool.

Dr. Cherry took Robinette et al's published data and divided the workers into three exposure levels. The above chart is the result of Dr. Cherry's analysis of the data set.

Robinette CD, Silverman C, Jablon S. Effects upon health of occupational exposure to microwave radiation (radar). *Am J Epidemiol* (1980); 112(1):39-53.

Cherry N. Health Effects in the vicinity of Radio/TV towers and mobile phone base stations. (2002): 1-40. <http://www.neilcherry.com/documents.php>

US Air Force (1970-1989)




US Air Force Workers with occupational exposure to microwave RF (1970-1989)

Y axis: Odds ratio for brain tumor

X axis: Exposure intensity score x months exposed)

Grayson JK. Radiation exposure, socioeconomic status, and brain tumor risk in the US Air Force: a nested case-control study. *Am J Epidemiol* (1996); 143(5):480-486.

Vatican Radio Tower (1987-1999).



Cumulative childhood leukaemia near the Vatican Radio Transmitters in Rome, 1987-1999.

Multiple powerful transmitters on site.

10 km radius around towers contains a population of >49,650 (1990 census).

exponential fitted trend line, $R^2=0.9756$, $p = 0.002$

Cherry N. Health Effects in the vicinity of Radio/TV towers and mobile phone base stations. (2002): 1-40.

<http://www.neilcherry.com/documents.php>

Michelozzi P, Capon A, Kirchmayer U et al. Adult and childhood leukemia near a high-power radio station in Rome, Italy. *Am J Epidemiol* (2002); 155(12):1096-1103.

Netanya, Israel (1997-1998)



New cell phone tower set up in city of Netanya, Israel, in July, 1996.

1500 watt, 850 MHz.


Power density in the whole exposed area was far below $0.53 \mu\text{w}/\text{cm}^2$.

This is 1000 times less than the FCC Guidelines of $600 \mu\text{W}/\text{cm}^2$ for 850 MHz exposure.

Comparison of cancer rates during the second year of exposure, in 677 long-term residents near the tower, compared to 1,222 matched controls living in another area of the city.

Wolf R, Wolf D. Increased Incidence of Cancer Near a Cell-Phone Transmitter Station. *International Journal of Cancer Prevention* (2004); 1(2):1-19.

Netanya, Israel - Relative Cancer Risk



Relative risk of cancer in residents near a new cell phone tower in Netanya, Israel, during the second year of exposure.


Overall risk of cancer in Area A was 4.15 times higher than in the town as a whole.

For men in area A, the cancer rate was 1.4 times higher.

For women in area A, the cancer rate was 10.5 times higher (p < 0.0001)

[the probability of this being a random finding is one hundredth of 1%

Naila, Germany (1999-2004)



Town of ~ 1100 residents.

Cell tower installed in 1993.

Medical of 1000 residents reviewed for the years 1994–2004.

Comparison of cancer incidents in residents living within 400 meters of the cell phone tower, compared to residents living farther away, and compared to the death rates for the province as a whole.

Eger H, Hagen K, Lucas B, Vogel P, Voit H. The Influence of Being Physically Near to a Cell Phone Transmission Mast on the Incidence of Cancer. *Umwelt·Medizin·Gesellschaft* (2004); 17(4):1-7.

Cancer Incidence in Naila (1999-2004)




Fig. 3 : Number of new cancer cases 1999 to 2004, adjusted for age and gender, calculated for the 5,000 patient years

Y axis: Cancer incidence 1994 – 2004 (new cases per 5000 patient years).

* Saarland = predicted rate based on the cancer registry for the federal state of Saarland.

** Naila = incidence for the town as a whole.


Inner area = residence within 400 meters of the tower.

Outer area = remainder of community.

In the inner area, the risk of cancer incidence was three times as high after five or more years of exposure.

In addition, the patients that live within 400 metres tend to develop the cancers at a younger age.

Hausmannstätten & Vasoldsberg, Austria (1984-1997)




NMT 450 cell tower, operational from 1984–1997.

Case/control study of cancer patients living within 1200 meter radius of the tower.

Oberfeld G. Environmental Epidemiological Study of Cancer Incidence in the Municipalities of Hausmannstätten & Vasoldsberg (Austria). Provincial Government of Styria, Department 8B, Provincial Public Health Office, Graz, Austria (2008):1-10. <http://www.emf-health.com/PDFreports/Austrianstudy.pdf>

Hausmannstätten & Vasoldsberg, Austria (1984-1997)



Odds ratio of cancer incidence — stratified by exposure levels (exterior to dwelling) in $\mu\text{W}/\text{m}^2$.

Note: FCC thermal safety guidelines ~ 6,000,000 $\mu\text{W}/\text{m}^2$)

In the highest exposure category:

Breast cancer risk was 23 times higher,

Brain cancer risk was 121 times higher.

Oberfeld G. Environmental Epidemiological Study of Cancer Incidence in the Municipalities of Hausmannstätten & Vasoldsberg (Austria). Provincial Government of Styria, Department 8B, Provincial Public Health Office, Graz, Austria (2008):1-10. <http://www.emf-health.com/PDFreports/Austrianstudy.pdf>

Belo Horizonte, Brazil (2011)




Belo Horizonte is the capital of Minas Gerais state in Brazil, population 2,258,096 in 2010.

Rated by the U.N. in 2007 as having the best quality of life in Latin America.

By 2006, 856 cell phone towers had been installed in the city.

Dode AC, Leao MM, Tejo Fde A et al. Mortality by neoplasia and cellular telephone base stations in the Belo Horizonte municipality, Minas Gerais state, Brazil. *Sci Total Environ* (2011); 409(19):3649-3665.



Environmental monitoring of RF power densities in the city was performed.


In 2003, the highest recorded power density in the city was $3.06 \mu\text{W}/\text{cm}^2$.

In 2008, the largest recorded power density was $40.78 \mu\text{W}/\text{cm}^2$, 13 times higher than in 2003.

$40 \mu\text{W}/\text{cm}^2$ is 15 times less than the FCC Exposure Guidelines.

Fig. 3. Horizontal and vertical radiation patterns per sector of BS site BH 20


From: Dode AC, Leao MM, Tejo Fde A et al. Mortality by neoplasia and cellular telephone base stations in the Belo Horizonte municipality, Minas Gerais state, Brazil. *Sci Total Environ* (2011); 409(19):3649-3665.



The authors used the Telecommunications National Agency database to map the locations of the 856 cell phone towers that existed in the city as of December 2006.

Fig. 8. Installed BSs in the Belo Horizonte municipality until 2006. Total amount = 856.

Dode AC, Leao MM, Tejo Fde A et al. Mortality by neoplasia and cellular telephone base stations in the Belo Horizonte municipality, Minas Gerais state, Brazil. *Sci Total Environ* (2011); 409(19):3649-3665.




They then cross-referenced health department records of death by neoplasia with census and demographic city population data to **locate the residence** of all individuals who had died of cancer in the city between 1996 and 2006.

Fig. 10. Sample of geocoded deaths and BS locations in downtown Belo Horizonte City located in Central-Southern region.

From: Dode AC, Leao MM, Tejo Fde A et al. Mortality by neoplasia and cellular telephone base stations in the Belo Horizonte municipality, Minas Gerais state, Brazil. *Sci Total Environ* (2011); 409(19):3649-3665.

Belo Horizonte, Brazil (2011)



Analysis of this data showed that the cancer death rate was significantly elevated at proximities closer than 500 meters to cell phone towers.

Fig. 15. Rate of mortality by neoplasia, according to the distance from the BS in Belo Horizonte municipality, from 1996 to 2006, and the null hypothesis (blue line).


Dode AC, Leao MM, Tejo Fde A et al. Mortality by neoplasia and cellular telephone base stations in the Belo Horizonte municipality, Minas Gerais state, Brazil. *Sci Total Environ* (2011); 409(19):3649-3665.

Effects of Microwave RF Exposure on Fertility



Human fertility is decreasing in the developed world.

Figure 2. Sperm concentration in 975 sperm donors recruited over 20 years



Sperm counts have been dropping worldwide for the last several decades.

(e.g. In New Zealand, 2.5% per year for the last 20 years).

Pesticides have been implicated.

Some evidence suggests that microwave RF exposure may also play a role.

Impaired Fertility in Fruit Flies



Insects are remarkably resistant to ionizing radiation and radioactivity.

They appear to be much more sensitive to the effects of microwave radio frequency exposures.


In a recent study, **fruit flies were exposed to $10 \mu\text{W}/\text{cm}^2$** of GSM 900 MHz or 1800 MHz digital RF.

This exposure level is 100 times lower than the FCC Guidelines of $1000 \mu\text{W}/\text{cm}^2$

Exposures were for one single exposure intervals per day for five days, ranging from 1 to 21 minutes per day.

Panagopoulos DJ, Margaritis LH. The effect of exposure duration on the biological activity of mobile telephony radiation. *Mutat Res* (2010); 699(1-2):17-22.

Impaired Fertility in Fruit Flies



0 = control group, with no exposure.

Even at one minute of exposure per day, a significant decrease in fertility is seen.

Fig. 2. Reproductive capacity (mean number of F1 pupae per maternal fly) of groups exposed to DCS 1800MHz radiation for different daily exposure durations (1, 6, 11, 16, and 21min) for five consecutive days, and of sham-exposed groups (no exposure).

Panagopoulos DJ, Margaritis LH. The effect of exposure duration on the biological activity of mobile telephony radiation. *Mutat Res* (2010); 699(1-2):17-22.


Impaired Fertility in Mammals



This is a Wistar rat.

A great deal of research has been done on the effects of microwave RF on laboratory animals.

Impaired Fertility in Female Mice



In one study, mice were kept in cages in a VHF/UHF antenna park in Thessaloniki, Greece.

Power densities ranged between 0.168 to $1.053 \mu\text{W}/\text{cm}^2$ [reported as 168 – 1053 nanowatts/ cm^2]


This is about 1000 times lower than the FCC Guidelines of $600\text{--}1000 \mu\text{W}/\text{cm}^2$

With repeated matings, litter size decreased, until by the 5th mating, all the dams were infertile.

This infertility was irreversible.

Magras IN, Xenos TD. RF radiation-induced changes in the prenatal development of mice. *Bioelectromagnetics* (1997); 18(6):455-461.

Impaired Fertility in Male Rats



Reduced sperm production in male Wistar rats exposed to 10 GHz microwave RF.

0.21 mW/cm² = **one fifth of the FCC Guidelines of 1 mW/cm²**

OTHER EFFECTS: Increases in reactive oxygen species, increased free radical formation, decreased activity of glutathione peroxidase and superoxide dismutase, DNA strand breakage, increased apoptosis (cell death) in sperm cells, distortion of sperm structure, reduced testosterone levels, shrinkage of seminiferous tubules and testicular size, decreased number and weight of progeny.


Kesari KK, Kumar S, Behari J. Effects of radiofrequency electromagnetic wave exposure from cellular phones on the reproductive pattern in male Wistar rats. *Biochem Biotechnol* (2011); 164(4):546-559.

Kesari KK, Kumar S, Behari J. Pathophysiology of microwave radiation: effect on rat brain. *Appl Biochem Biotechnol* (2012); 166(2):379-388.

Kumar S, Kesari KK, Behari J. Influence of microwave exposure on fertility of male rats. *Fertil Steril* (2011); 95(4):1500-1502.

Kumar S, Behari J, Sisodia R. Influence of electromagnetic fields on reproductive system of male rats. *Int J Radiat Biol* (2012); epub Nov 13:1-8

WiFi Exposure Damages Sperm With Oxidant Stress.



The rats were exposed to a Standard WiFi gateway, 24 hours a day for 20 days.

HSCORE = histological staining in testes for 8-OH-20-dG
[8-hydroxy-20-deoxyguanosine, **byproduct of DNA damage**]

Serum 8-OH-20-dG (ng/ml) [byproduct of DNA damage]

TBS = testicular biopsy score

9 = Much spermatogenesis, but germinal epithelium disorganized with marked sloughing or obliteration of lumen

GPX = glutathione peroxidase, an antioxidant (consumed by oxidative stress in exposed rats).

Atasoy HI, Gunal MY, Atasoy P, Elgun S, Bugdayci G. Immunohistopathologic demonstration of deleterious effects on growing rat testes of radiofrequency waves emitted from conventional Wi-Fi devices. *J Pediatr Urol* (2012); March 30.

Impaired Fertility in Birds




In Valladolid, Spain, a study compared the productivity of storks nesting closer and farther from a cell phone tower site.

30 nests within 200 meters of the antennae, were compared with 30 nests greater than 300 meters from the antennae

Balmori A. Possible Effects of Electromagnetic Fields from Phone Masts on a Population of White Stork. *Electromagn Biol Med* (2005); 24(2):109-119.

Impaired Fertility in Birds



Productivity was significantly reduced in birds in the high exposure group.

Average electric field intensity on nests within 200m = 2.36 ± 0.82 V/m ($\sim 1.48 \mu\text{W}/\text{cm}^2$)

This is more than 400 times less than the FCC Guidelines of 600–1000 $\mu\text{W}/\text{cm}^2$

Average electric field intensity on nests further than 300m = 0.53 ± 0.82 V/m ($\sim 0.07 \mu\text{W}/\text{cm}^2$).

Impaired Fertility in Amphibians




Eggs and tadpoles of the European common frog (*Rana temporaria*) were exposed to RF/EFM from several cell towers located at a distance of 140 meters.

Duration of exposure was 2 months (from egg phase to advanced tadpole stage).

Control groups were placed in same conditions, but contained in a faraday cage that shielded the eggs from RF exposure.

Balmori A. Mobile phone mast effects on common frog (*Rana temporaria*) tadpoles: the city turned into a laboratory. *Electromagn Biol Med* (2010a); 29(1-2):31-35.

Impaired Fertility in Amphibians




Exposure intensity 1.8 to 3.5 V/m ($\sim 0.8\text{--}3.2 \mu\text{W}/\text{cm}^2$).

This is 200 times less than the FCC Guidelines of $600\text{--}1000 \mu\text{W}/\text{cm}^2$

[In the exposed group ($n = 70$), low coordination of movements and asynchronous growth was observed in living specimens, resulting in both big and small tadpoles. In the control group ($n = 70$), growth was normal.]

Balmori A. Mobile phone mast effects on common frog (*Rana temporaria*) tadpoles: the city turned into a laboratory. *Electromagn Biol Med* (2010a); 29(1-2):31-35.

Cell Phone Transmissions Decrease Sperm Motility in Vitro



Samples of human sperm received 5 minutes exposure, 10 cm from a transmitting GSM 900 MHz cell phone.


Average power density of exposure: $20 \mu\text{W}/\text{cm}^2$

This is 30 times less than the FCC Exposure Guideline of $600 \mu\text{W}/\text{cm}^2$

(Y axis = values in %)

Erogul O, Oztas E, Yildirim I et al. Effects of electromagnetic radiation from a cellular phone on human sperm motility: an in vitro study. *Arch Med Res* (2006); 37(7):840-843.


Cell Phone Use Decreases Sperm Motility in Vivo



Semen analysis performed on 371 men at a university clinic.
Health questionnaire included query of cell phone use habits.
(Y axis = values in %)

Fejes I, Zavaczki Z, Szollosi J et al. Is there a relationship between cell phone use and semen quality? *Arch Androl* (2005); 51(5):385-393.

Cell Phone Use Degrades Sperm Quality in Vivo



Three hundred sixty-one men undergoing infertility evaluation were divided into four groups according to their active cell phone use:


group A: no use; group B: <2 h/day; group C: 2-4 h/day; and group D: >4 h/day.

With greater than two hours a day of reported talk time, significant reduction in **sperm count, motility, viability, and % normal morphology** were observed.

[One can assume that with texting rather than talking, the data might be even worse . . . as the phone antenna will be closer to the testes.]

Agarwal A, Deepinder F, Sharma RK, Ranga G, Li J. Effect of cell phone usage on semen analysis in men attending infertility clinic: an observational study. *Fertil Steril* (2008); 89(1):124-128.

Isothermal Exposure to 1.8 GHz RF Damages Sperm




Sperm exposed for 16 hours in vitro to 1.8 GHz (SAR = 27.5 W/kg) @ 21°C (isothermal conditions).

Sperm damage correlates with increased free radical (ROS) production.

Values in %.

De Iuliis GN, Newey RJ, King BV, Aitken RJ. Mobile phone radiation induces reactive oxygen species production and DNA damage in human spermatozoa in vitro. *PLoS One* (2009); 4(7):e6446 (1-9).

1.8 GHz RF Degrades Sperm Quality In Vitro



1.8 GHz RF at various intensities for 16 hours @ 21°C

This is an isothermal exposure


Sperm vitality and motility are significantly detruded at SAR = 1 W/kg and above

Figure 2. RF-EMR exposure reduces motility and vitality of human spermatozoa, in an SAR dependent manner. Percoll-purified spermatozoa (5×10^6 cells) were suspended in 1 ml BWW in a 35 mm Petri dish and placed within the waveguide while control cells (closed circles) were placed outside the waveguide. Cells in the waveguide were exposed to 1.8 GHz RF-EMR at SAR levels of 0.4, 1.0 2.8 4.3 10.1 and 27.5 W/kg (open circles) for 16 h at 21uC. Both vitality and motility were reduced in a dose dependent manner.

A, Vitality was significantly reduced at a SAR of 1.0 W/kg from 89% \pm 3% to 65% \pm 1% (**p,0.01).

B, Motility was also significantly reduced at a SAR of 1.0 W/kg from 86% \pm 2% to 68% \pm 2% (*p,0.05). All results are based on 4 independent samples.

ROS Production – RF versus Thermal




A. ROS generation (DHE response) was significantly increased from control levels after exposure to 1.0 W/kg (*p, 0.05) and above (***p, 0.001).

C. In order to control for thermal effects, the impact of temperature of cellular ROS generation was monitored; a significant increase in ROS generation was observed as temperatures rose above 40°C (p, 0.001).

Figure 3. RF-EMR induces ROS generation in human spermatozoa, in an SAR-dependent manner unrelated to thermal effects.

De Iuliis GN, Newey RJ, King BV, Aitken RJ. Mobile phone radiation induces reactive oxygen species production and DNA damage in human spermatozoa in vitro. *PLoS One* (2009); 4(7):e6446 (1-9).

Oxidative Damage To Sperm DNA From 1.8 GHz RF Exposure



1.8 GHz RF x 16 hours @ 21°C isothermal.

A) As the power levels were increased, the amount of oxidative DNA damage expressed also increased.


A significant amount of oxidative DNA damage was observed in cells exposed to 2.8 W/kg (*p,0.05) RF-EMR and above (**p,0.01; ***p,0.001).

B) The levels of 8-OH-dG expression were positively correlated with the levels of ROS generation by the mitochondria ($R^2 = 0.727$).

Figure 4. RF-EMR induces oxidative DNA damage in human spermatozoa.

De Iuliis GN, Newey RJ, King BV, Aitken RJ. Mobile phone radiation induces reactive oxygen species production and DNA damage in human spermatozoa in vitro. *PLoS One* (2009); 4(7):e6446 (1-9).

RF Damages Sperm by Increasing Oxidative Stress



A) Significant levels of DNA fragmentation was observed in exposed spermatozoa at 2.8 W/kg (*p,0.05) and above (**p,0.001).

B) DNA fragmentation was positively correlated with ROS production by the mitochondria as monitored by MSR. ($R^2 = 0.861$).

Figure 5. RF-EMR induces DNA fragmentation in human spermatozoa.

De Iuliis GN, Newey RJ, King BV, Aitken RJ. Mobile phone radiation induces reactive oxygen species production and DNA damage in human spermatozoa in vitro. *PLoS One* (2009); 4(7):e6446 (1-9).

Sperm Damage From Laptop WiFi




Motile spermatozoa in semen were incubated at room temperature,
3 cm below laptop computer (e.g. lap distance)
4 hours of exposure.

Control incubated in similar conditions, without presence of the computer.

Avendano C, Mata A, Sanchez Sarmiento CA, Doncel GF. Use of laptop computers connected to internet through Wi-Fi decreases human sperm motility and increases sperm DNA fragmentation. *Fertil Steril* (2012); 97(1):39-45.

Sperm Damage From Laptop WiFi



Power density ranged 0.45 to 1.05 $\mu\text{W}/\text{cm}^2$

[This is roughly **1000 times less** than the FCC exposure limit of 1000 $\mu\text{W}/\text{cm}^2$]

Avendano C, Mata A, Sanchez Sarmiento CA, Doncel GF. Use of laptop computers connected to internet through Wi-Fi decreases human sperm motility and increases sperm DNA fragmentation. *Fertil Steril* (2012); 97(1):39-45.

Cell Phones and Risk of Brain Tumor



THEME: CANCER

Brain tumour risk in relation to mobile telephone use: results of the INTERPHONE international case–control study

The INTERPHONE Study Group*

Corresponding author. Elisabeth Cardis; CREAL, Doctor Aiguader 88, 08003 Barcelona, Spain. E-mail: ecardis@creal.cat

*List of members of this study group is available in the Appendix.

Accepted 8 March 2010

Background The rapid increase in mobile telephone use has generated concern about possible health risks related to radiofrequency electromagnetic fields from this technology.

Methods An interview-based case–control study with 2708 glioma and 2409 meningioma cases and matched controls was conducted in 13 countries using a common protocol.

Large case–control study at multiple centers.


Funded in significant part by the telecommunications industry.

INTERPHONE Study 2010

Conclusions Overall, no increase in risk of glioma or meningioma was observed with use of mobile phones.

From the authors' published conclusions.

Odds Ratio for Meningioma and Glioma



In the 2010 Interphone Study combined analysis of data for all levels of exposure found that “regular cell phone users” were less likely to have brain tumors than non-users.

This is what was reported in the media about this study.

A reduced odds ratio (OR) related to ever having been a regular mobile phone user was seen for glioma [OR 0.81; 95% confidence interval (CI) 0.70–0.94] and meningioma (OR 0.79; 95% CI 0.68–0.91)

INTERPHONE Study Group. Brain tumour risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Int J Epidemiol* (2010); 39(3):675-694.


INTERPHONE Study 2010 – Definition of “Risk”

if the subject had ever been a regular user of a mobile phone (had an average of at least one call per week for a period of ≥ 6 months).²⁶

However, “regular use” was defined as a minimum of one call for week for at least 6 months.

In otherwords, anyone who had made at least 26 cell phone calls in their lifetime was categorized as a “regular user” and placed in the risk group.

Odds Ratio for Meningioma and Glioma with Cell Phone Use



Cumulative call time without hands-free devices, **divided into deciles.**

Blue line is Odds Ratio of 1.0 (equal to control group).


Half of the subjects in the study had less than 115 hours of lifetime exposure.

Note that a significantly higher risk for glioma was seen with more than 1640 hours of exposure.

OR for glioma = 1.40 [95% CI = (1.03-1.89)]

From Table 2: INTERPHONE Study Group. Brain tumour risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Int J Epidemiol* (2010); 39(3):675-694.

Odds Ratio for Meningioma with Cell Phone Use



Charted data from the Interphone study for risk of meningioma.

With ≥ 1640 hrs exposure in 1 – 4 years, OR = 4.80 [95% CI = (1.49–15.4)]


1640 hours in 4 years = 7.9 hrs/wk (range in cohort was 8 – 30 hrs/wk)

1640 hours in 10+ years ~ > 3 hrs/wk

(range in 1–4 years cohort was 8 – 30 hrs/wk, which the authors discounted as “implausible values of use” in their summary of results)

From Table 3: INTERPHONE Study Group. Brain tumour risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Int J Epidemiol* (2010); 39(3):675-694.

Odds Ratio for Glioma with Cell Phone Use



Charted data from INTERPHONE study group, glioma risk.

Stratified by cumulative call time (without hands/free devices).

Also stratified by years of use.


With ≥ 1640 hrs exposure in 1 - 4 years, OR for Glioma = 3.77 [95% CI = (1.25-11.4)]

1640 hours in 4 years = 7.9 hrs/wk (range in cohort was 8 - 30 hrs/wk)

The authors rejected their own findings on glioma, stating that this level of reported cell phone use (8 to 30 hours per week) was “implausible”.

From Table 3: INTERPHONE Study Group. Brain tumour risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Int J Epidemiol* (2010); 39(3):675-694.

Odds Ratio for Glioma with Cell Phone Use



Glioma risk by **location in the brain.**


Also stratified by **cumulative call time.**

Temporal lobe (With ≥ 1640 hrs exposure, OR = 1.87 [95% CI = (1.09-3.22)]

From Table 4: INTERPHONE Study Group. Brain tumour risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Int J Epidemiol* (2010); 39(3):675-694.

Interestingly, the study did report its statistics stratified by total time of reported use, and the top decile (greater than 1640 hours use over a ten year interval, averaging out as greater than 3 hours a week) had an increased risk of certain tumors. Individuals who accrued that greater than 1650 hours of use over a 1 to 4 year interval (ranging from 8 to over 30 hours a week) had a markedly higher odds ratio of meningioma (OR 4.80) or glioma (OR 3.27).

Odds Ratio for Glioma with Cell Phone Use




Glioma risk by **side of head they habitually held the phone.**

Also stratified by **cumulative call time.**


(With ≥ 1640 hrs exposure, Ipsilateral OR = 1.96 [95% CI = (1.22-3.22)])

Odds Ratio for Glioma (by Years of Use)



Stratified by years of exposure.

Odds Ratio for Glioma (by Years of Use)



Glioma risk by **side of head they habitually held the phone.**

Also stratified by **years of exposure.**

INTERPHONE Study 2011

Acoustic neuroma risk in relation to mobile telephone use: Results of the INTERPHONE international case-control study

The INTERPHONE Study Group^{*,1}

ARTICLE INFO

Article history:

Received 13 April 2011

Received in revised form 9 May 2011

Accepted 10 May 2011

Available online 23 August 2011

Keywords:

Acoustic neuroma

Vestibular schwannoma

Brain tumour

Mobile phones

Radiofrequency electromagnetic fields

Epidemiology

ABSTRACT

Background: The rapid increase in mobile telephone use has generated concern about possible health risks of radiofrequency electromagnetic fields from these devices. *Methods:* A case-control study of 1105 patients with newly diagnosed acoustic neuroma (vestibular schwannoma) and 2145 controls was conducted in 13 countries using a common protocol. Past mobile phone use was assessed by personal interview. In the primary analysis, exposure time was censored at one year before the reference date (date of diagnosis for cases and date of diagnosis of the matched case for controls); analyses censoring exposure at five years before the reference date were also done to allow for a possible longer latent period. *Results:* The odds ratio (OR) of acoustic neuroma with ever having been a regular mobile phone user was 0.85 (95% confidence interval 0.69–1.04). The OR for ≥ 10 years after first regular mobile phone use was 0.76 (0.52–1.11). There was no trend of increasing ORs with increasing cumulative call time or cumulative number of calls, with the lowest OR (0.48 (0.30–0.78)) observed in the 9th decile of cumulative call time. In the 10th decile (≥ 1640 h) of cumulative call time, the OR was 1.32 (0.88–1.97); there were, however, implausible values of reported use in those with ≥ 1640 h of accumulated mobile phone use. With censoring at 5 years before the reference date the OR for ≥ 10 years after first regular mobile phone use was 0.83 (0.58–1.19) and for ≥ 1640 h of cumulative call time it was 2.79 (1.51–5.16), but again with no trend in the lower nine deciles and with the lowest OR in the 9th decile. In general, ORs were not greater in subjects who reported usual phone use on the same side of the head as their tumour than in those who reported it on the opposite side, but it was greater in those in the 10th decile of cumulative hours of use. *Conclusions:* There was no increase in risk of acoustic neuroma with ever regular use of a mobile phone or for users who began regular use 10 years or more before the reference date. Elevated odds ratios observed at the highest level of cumulative call time could be due to chance, reporting bias or a causal effect. As acoustic neuroma is usually a slowly growing tumour, the interval between introduction of mobile phones and occurrence of the tumour might have been too short to observe an effect, if there is one.


© 2011 Elsevier Ltd. All rights reserved.

2011 INTERPHONE study of acoustic neuroma

Funded in significant part by the telecommunications industry.

Cardis E, Schüz J. Acoustic neuroma risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Cancer Epidemiol* (2011); 35(5):453-464.

Odds Ratio for Acoustic Neuroma with Cell Phone Use



From 2011 INTERPHONE study of acoustic neuroma, Table 2.

This study was also funded in major part by the telecommunications industry.


In their conclusion, the authors stated that this data showed **“no trend of increasing risk with increasing cumulative call time”**

They discounted their findings for the highest decile of exposure.

But with ≥ 1640 hrs exposure in 1 - 5 years of exposure, **OR = 2.79 [95% CI = (1.51-5.16)]**

From Table 2: Cardis E, Schüz J. Acoustic neuroma risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Cancer Epidemiol* (2011); 35(5):453-464.

Acoustic Neuroma Risk (< 1 year of cell phone use)




In the high use group, risk of acoustic neuroma was significantly higher on the side of the head where the subject habitually held the cell phone.

1640 hours in 1 year = 4.5 hours a day = 31.5 hours/week

With ≥ 1640 hrs exposure, ipsilateral tumor Odds Ratio = 2.33 [95% CI = (1.23–4.40)]

From Table 3: Cardis E, Schüz J. Acoustic neuroma risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Cancer Epidemiol* (2011); 35(5):453-464.

Acoustic Neuroma Risk (< 5 years of cell phone use)




Risk increased with increased years of exposure.

1640 hours in 5 years = 0.9 hours a day = 6.3 hours/week

With ≥ 1640 hrs exposure, ipsilateral tumor Odds Ratio = 3.53 [95% CI = (1.59–7.82)]

From Table 3: Cardis E, Schüz J. Acoustic neuroma risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Cancer Epidemiol* (2011); 35(5):453-464.

Acoustic Neuroma Risk (≥ 10 years of cell phone use)




Higher risk with ten or more years of exposure.

1640 hours in 10 years = less than half an hour a day. = 3.2 hours/week = 0.45 hours a day

With ≥ 1640 hrs exposure, ipsilateral tumor Odds Ratio = 3.74 [95% CI = (1.58–8.83)]

From Table 4: Cardis E, Schüz J. Acoustic neuroma risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Cancer Epidemiol* (2011); 35(5):453-464.

Acoustic Neuroma Risk with 1 to 4 Years of Cell Phone Use




2011 INTERPHONE study of acoustic neuroma

From Table 4: Short, medium, long-term accumulation of >1640 hours.

Cardis E, Schüz J. Acoustic neuroma risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Cancer Epidemiol* (2011); 35(5):453-464.

Acoustic Neuroma Risk with 5 to 9 Years of Cell Phone Use




2011 INTERPHONE study of acoustic neuroma

From Table 4: Short, medium, long-term accumulation of >1640 hours.

Cardis E, Schüz J. Acoustic neuroma risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Cancer Epidemiol* (2011); 35(5):453-464.

Acoustic Neuroma Risk with 10 + Years of Cell Phone Use



2011 INTERPHONE study of acoustic neuroma

From Table 4: Short, medium, long-term accumulation of ≥ 1640 hours.

Cardis E, Schüz J. Acoustic neuroma risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Cancer Epidemiol* (2011); 35(5):453-464.



Risk of brain tumours in relation to estimated RF dose from mobile phones: results from five Interphone countries

E Cardis,¹ B K Armstrong,² J D Bowman,³ G G Giles,^{4,5} M Hours,⁶ D Krewski,⁷ M McBride,⁸ M E Parent,⁹ S Sadetzki,^{10,11} A Woodward,¹² J Brown,² A Chetrit,¹⁰ J Figuerola,¹ C Hoffmann,^{11,13} A Jarus-Hakak,¹⁰ L Montestrucq,⁶ L Nadon,⁹ L Richardson,¹⁴ R Villegas,¹ M Vrijheid¹

For numbered affiliations see end of article.

Correspondence to

Professor E Cardis, Centre for Research in Environmental Epidemiology (CREAL), Hospital del Mar Research Institute (IMIM), CIBER Epidemiología y Salud Pública (CIBERESP), Doctor Aiguader 88, 08003 Barcelona, Spain; ecardis@creal.cat

Accepted 20 May 2011
Published Online First
9 June 2011

ABSTRACT

Objectives The objective of this study was to examine the associations of brain tumours with radio frequency (RF) fields from mobile phones.

Methods Patients with brain tumour from the Australian, Canadian, French, Israeli and New Zealand components of the Interphone Study, whose tumours were localised by neuroradiologists, were analysed. Controls were matched on age, sex and region and allocated the 'tumour location' of their matched case. Analyses included 553 glioma and 676 meningioma cases and 1762 and 1911 controls, respectively. RF dose was estimated as total cumulative specific energy (TCSE; J/kg) absorbed at the tumour's estimated centre taking into account multiple RF exposure determinants.

Results ORs with ever having been a regular mobile phone user were 0.93 (95% CI 0.73 to 1.18) for glioma and 0.80 (95% CI 0.66 to 0.96) for meningioma. ORs for glioma were below 1 in the first four quintiles of TCSE

What this paper adds

- ▶ Previous epidemiological studies of mobile phone use and brain cancer risk have used information on mobile phone use as a proxy measure of exposure to radio frequency fields from mobile phones.
- ▶ Most studies have not observed increased ORs in relation to ever having been a mobile phone user. There were suggestions, however, of an increased risk of glioma in long-term and heavy users, though biases and errors prevent a causal interpretation.
- ▶ The relationship between radio frequency energy absorbed at the tumour location and mobile phone use history is complex. In addition to amount of use, it depends on phone type, network properties, conditions of use and

Interphone latest study 2011

Cardis E, Armstrong BK, Bowman JD et al. Risk of brain tumours in relation to estimated RF dose from mobile phones: results from five Interphone countries. *Occup Environ Med* (2011); 68(9):631-640.

INTERPHONE Study 2011

“Our results suggest that there may be an increase in risk of glioma in the most exposed area of the brain among long-term and heavy users of mobile phones. These results are uncertain (in light of the uncertainties associated with tumour centre localisation, radio frequency dose estimation and sample size) and require replication before they can be taken to indicate a cause effect relationship.”

From the conclusions of the 2001 INTERPHONE study (industry-financed).

Study authors finally admitted that their data showed increase risk of glioma, but said that this finding required replication before being taken as a cause and effect relationship.

This despite the fact that this finding was already a replication of their previously published data, and had also been confirmed several times in the published data of the Hardell group in Sweden.

BBC NEWS

HEALTH

20 October 2011 Last updated at 21:11 ET

Mobile phone brain cancer link rejected

By Nick Triggle

Health correspondent, BBC News

Further research has been published suggesting there is no link between mobile phones and brain cancer.

The risk mobiles present has been much debated over the past 20 years as use of the phones has soared.

Danish study: Proclaimed as evidence that cell phones are safe.

420,095 subscribers in the cohort — who had subscriptions by 1994/95.


Exposure is judged by presence of a cell phone contract, no record of actual usage.

200,507 corporate users excluded — **and placed in the control group.**

2550 juveniles excluded — **and placed in the control group.**

Frei P, Poulsen AH, Johansen C, Olsen JH, Steding-Jessen M, Schuz J. Use of mobile phones and risk of brain tumours: update of Danish cohort study. *BMJ* (2011); 343(d6387).

Odds Ratio for Brain Tumor (7+ years of cell phone use)



Stratified by Total Cumulative Energy Exposure (joules/kilogram)

From Table 3: Cardis E, Armstrong BK, Bowman JD et al. Risk of brain tumours in relation to estimated RF dose from mobile phones: results from five Interphone countries. *Occup Environ Med* (2011); 68(9):631-640.

Danish Study 2011

Danish study: Proclaimed as evidence that cell phones are safe.

420,095 subscribers in the cohort — who had subscriptions by 1994/95. Exposure is judged by presence of a cell phone contract, no record of actual usage.

200,507 corporate users excluded — and placed in the control group.

2550 juveniles excluded — and placed in the control group.

Half the subjects in the 2009 Johansen et al study had less than two years of cell phone use.

Second publication: Schuz 2006

Only 61% of subscribers reported making or receiving at least 1 call a week in prior six months

All users who began subscription after 1995 were put in the “unexposed” reference population.

Third publication Shuz et al 2011

Same study group

Control group 2.9 million Danes

Fourth publication: Frei et al 2011 BMJ

~ 42% of initial cohort excluded (*and placed in the control group*).

Also in the control group — the 85% of Danes that got a cell phone contract between 1995 and 2004.

“Number of subscription years” is used as a surrogate for actual hours of usage.

18–29 year old excluded.

Cohort established by grants from Danish telcom companies. Sources of funding of the International Epidemiology Institute (Rockville, MD, USA) have never been declared.

In this study, the control group was contaminated with so many cell phone users that the results of the study were essentially meaningless. Two reviews stating this fact were published in the same issue of BMJ in 2011, along with the Frie study.

Frei P, Poulsen AH, Johansen C, Olsen JH, Steding-Jessen M, Schuz J. Use of mobile phones and risk of brain tumours: update of Danish cohort study. *BMJ* (2011); 343(d6387).

Philips A, Lamburn G. Updated study contains poor science and should be disregarded. *BMJ* (2011); 343(d7899; author reply d7912).

Soderqvist F, Carlberg M, Hardell L. Review of four publications on the Danish cohort study on mobile phone subscribers and risk of brain tumors. *Rev Environ Health* (2012); 27(1):51-58.

Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects

LENNART HARDELL¹, MICHAEL CARLBERG¹ and KJELL HANSSON MILD²

¹Department of Oncology, University Hospital, SE-701 85 Örebro;

²Department of Radiation Physics, Umeå University, SE-901 87 Umeå, Sweden

Received December 6, 2010; Accepted January 20, 2011

DOI: 10.3892/ijo.2011.947

Abstract. We studied the association between use of mobile and cordless phones and malignant brain tumours. Pooled analysis was performed of two case-control studies on patients with malignant brain tumours diagnosed during 1997-2003 and matched controls alive at the time of study inclusion and one case-control study on deceased patients and controls diagnosed during the same time period. Cases and controls or relatives to deceased subjects were interviewed using a structured questionnaire. Replies were obtained for 1,251 (85%) cases and 2,438 (84%) controls. The risk increased with

emissions from wireless devices such as mobile phone base stations, broadcast transmission towers, pagers and personal digital assistants, wireless networks and other sources of RF radiation (1).


The brain is the target organ of the body with highest near field exposure to microwaves during use of a handheld wireless phone. Thus, fear of an increased risk for brain tumours from RF fields emitted from mobile phones has dominated the debate the last decade. Of equal importance is use of the desktop cordless phones.

The most reliable research on the tumor risks of cell phones has been performed by the Hardell group in Sweden.

This group does not receive funding from the cell phone industry.

This is the only group that has controlled for use of in-home cordless phones as well as cell phones [which makes their data more reliable].

Tumor Risk by Years of Use – Glioma




Orient to the bar graph.

Hardell group -- current summary

From Table IV: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Tumor Risk by Years of Use – Astrocytoma



Hardell group -- current summary

From Table IV: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Risk of All Brain Tumors (Usage Below Median)



Hardell group -- current summary

From Table IV: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Risk of All Brain Tumors (Usage Above Median)



Hardell group -- current summary

From Table IV: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Risk of Glioma (Usage Below Median)



Hardell group -- current summary

From Table IV: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Risk of Glioma (Usage Above Median)



Hardell group -- current summary

From Table IV: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Risk of Astrocytoma (Usage Below Median)



Hardell group -- current summary

From Table IV: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Risk of Astrocytoma (Usage Above Median)



Hardell group -- current summary

From Table IV: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Risk of Oligodendroglioma (Usage Below Median)



Hardell group -- current summary

From Table IV: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Risk of Oligodendroglioma (Usage Above Median)



Hardell group -- current summary

From Table IV: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Risk of Tumor by Location – All Tumor Types



Hardell group -- current summary

From Table 3: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of two case-control studies on use of cellular and cordless telephones and the risk for malignant brain tumours diagnosed in 1997-2003. *Int Arch Occup Environ Health* (2006b); 79(8):630-639.


Risk of Tumor by Location – High Grade Astrocytoma



Hardell group -- current summary

From Table 3: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of two case-control studies on use of cellular and cordless telephones and the risk for malignant brain tumours diagnosed in 1997-2003. *Int Arch Occup Environ Health* (2006b); 79(8):630-639.


Risk of Tumor by Location – Low Grade Astrocytoma



Hardell group -- current summary

From Table 3: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of two case-control studies on use of cellular and cordless telephones and the risk for malignant brain tumours diagnosed in 1997-2003. *Int Arch Occup Environ Health* (2006b); 79(8):630-639.


Risk of Tumor by Location – Other Malignant Tumors



Hardell group -- current summary

From Table 3: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of two case-control studies on use of cellular and cordless telephones and the risk for malignant brain tumours diagnosed in 1997-2003. *Int Arch Occup Environ Health* (2006b); 79(8):630-639.


Tumor Risk by Cumulative Hours of Use – Any Brain Cancer



Hardell group -- current summary

From Table III: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Tumor Risk by Cumulative Hours of Use – Glioma



Hardell group -- current summary

From Table III: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Tumor Risk by Cumulative Hours of Use – Astrocytoma



Hardell group -- current summary

From Table III: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Tumor Risk by Cumulative Hours of Use – Oligodendroglioma



Hardell group -- current summary

From Table III: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Tumor Risk by Cumulative Hours of Use – Other/Mixed Glioma



Hardell group -- current summary

From Table III: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Tumor Risk by Cumulative Hours of Use – Other Brain Malignancy



Hardell group -- current summary

From Table III: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.


Tumor Risk by Age of First Use – Any Brain Cancer



Hardell group -- current summary

From Table V: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.

Tumor Risk by Age of First Use – Astrocytoma



Hardell group -- current summary

From Table V: Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* (2011b); 38(5):1465-1474.

Mobile phone use and brain tumours in the CERENAT case-control study

Gaëlle Coureau,^{1,2,3} Ghislaine Bouvier,^{1,2} Pierre Lebailly,^{4,5,6} Pascale Fabbro-Peray,^{7,8} Anne Gruber,¹ Karen Leffondre,² Jean-Sebastien Guillamo,⁹ Hugues Loiseau,¹⁰ Simone Mathoulin-Pélissier,² Roger Salamon,^{2,3} Isabelle Baldi^{1,2,11}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/oemed-2013-101754>).

For numbered affiliations see end of article.

Correspondence to

Dr Gaëlle Coureau, Université Bordeaux Segalen, ISPED, Equipe Santé Travail Environnement, 146 rue Léo Saignat, 33076 Bordeaux, Cedex, France; gaelle.coureau@isped.u-bordeaux2.fr

Received 23 July 2013
Revised 7 April 2014
Accepted 15 April 2014
Published Online First
9 May 2014

ABSTRACT

The carcinogenic effect of radiofrequency electromagnetic fields in humans remains controversial. However, it has been suggested that they could be involved in the aetiology of some types of brain tumours.

Objectives The objective was to analyse the association between mobile phone exposure and primary central nervous system tumours (gliomas and meningiomas) in adults.

Methods CERENAT is a multicenter case-control study carried out in four areas in France in 2004–2006. Data about mobile phone use were collected through a detailed questionnaire delivered in a face-to-face manner. Conditional logistic regression for matched sets was used to estimate adjusted ORs and 95% CIs.

Results A total of 253 gliomas, 194 meningiomas and 892 matched controls selected from the local electoral rolls were analysed. No association with brain tumours was observed when comparing regular mobile phone users with non-users (OR=1.24; 95% CI 0.86 to 1.77 for gliomas, OR=0.90; 95% CI 0.61 to 1.34 for meningiomas). However, the positive association was statistically significant in the heaviest users when


What this paper adds

- The potential association between mobile phone use and brain tumour remains controversial, and original data have mostly been provided by studies performed in Sweden and the international Interphone study.
- Some studies suggest that long-term (over 10 years) mobile phone use increases the risk of gliomas, and especially of those with temporal location.
- This analysis highlights a positive association between heavy use of mobile phone and brain tumour, considering life-long cumulative duration and number of calls.
- Risks were higher for gliomas, temporal tumours, occupational and urban mobile phone use.
- This study provides additional data supporting a possible association between heavy mobile phone use and brain tumours.

The increased risk of brain tumor with cell phone has been confirmed in another study from France, published in May 2014

Coureau G, et al. *Occup Environ Med* 2014;71:514–522. doi:10.1136/oemed-2013-101754

Risk of glioma with cell phone use.



What is the long-term outlook for WiFi in the Classroom?



If wifi is installed in the classroom, at least 3–5% of the exposed individuals are likely to develop some symptoms.

Increased levels of headache, concentration difficulty, memory problems, tinnitus, irritability, and hyperactivity or depression could be expected to occur in sensitive individuals. This would include both students and staff. Is the school system prepared to address the needs of these individuals?

The evidence on acute effects and on long-term degenerative changes produced by chronic RF exposure will continue to build over the next few years.

Public concern is likely to continue to grow over time, especially as regards the exposure of children in an environment where attendance is required by law.

These problems and concerns will have legal repercussions, down the line.

Constant wifi transmissions should be avoided.



Schools that are using wifi to communicate with computers and tablets should avoid curriculum that requires constant interaction across the network, and choose curricular elements that can be used in a “stand-alone” format once they are loaded onto the tablets.

Wifi transmitters should be turned off when they are not being actively used, to eliminate their constant transmission. Laptops and tablets should be placed in airplane mode when active communication across the network is not required.

These measures could reduce the exposure to wifi transmissions significantly, and might help to reduce (to some extent) the amount of pathology that will be produced by the use of this technology.

'LiFi' uses LEDs for blazing fast data transfer

Nick Barber
@nickjb

Nov 7, 2013 4:45 AM

A new technology called LiFi can transfer data using LED lights. In this video, we'll watch a demo of a LiFi system made with off-the-shelf-parts, as it streams a vide.




The development of “LiFi” may offer a more convenient option in the future. Lifi will require line of sight between transmitter and receiver on the mobile device. But it will have much more bandwidth than wifi, so the market will probably go that way. This technology will probably start to appear in the marketplace in 2015.

Even if the school systems install WiFi now, they are likely to retreat to cabled connections or LiFi in the future, for medical and legal reasons.

That will be an expensive retreat.

A cabled connection would be a better choice.



Cabled connections are less convenient logistically, but avoid the concerns we've addressed here.

Tablet or eReader devices that can connect to a computer via USB cable would be preferable to devices that can only communicate by wifi.

Basic Precepts for Residential RF Exposures



Proposed to the Eugene Water and Electric Board by their medical advisory group (Eugene, Oregon, July 2013).

RECOMMENDATIONS r.e. SMART METERS:

EWEB should adopt a policy of minimizing their RF footprint in the residential community.

I'm through describing the problem now. If you'd like, I can briefly review our group's recommendations to the Board.

If EWEB chooses to go ahead with RF smart meters, the solution is to use these tools in a safer way.

Design the system to truly minimize the amount of RF signal that is being produced.

Most engineering choices in AMI have not been designed with these goals in mind.

The industry has devoted more energy to denying risk than it has to designing systems that are "smart" enough to minimize that risk.

But our group thinks that it could be possible to take a prudent approach and still get the job done.

What can EWEB do to actually "work the problem"? Let's summarize the situation:

The Problem:

Excessive RF exposure can cause acute symptoms in many people.

As we've shown, some people get acute symptoms from excessive RF exposures (headaches, insomnia, fatigue, vertigo, tinnitus, other symptoms of EHS).

People like this exist in Eugene.

How many people might this be in our town?

Based on existing research, a conservative guess might be 3 to 5% of the population.

3% of 150,000 people is 4,500 people.

The Problem:

Excessive RF exposure can also cause severe chronic problems.

Chronic RF exposure increases oxidative stress in living systems.

Increased cancer and male infertility are only two of many problems potentially caused by chronic elevation of oxidative stress.

Allowing an
“opt out” is not
a practical
solution.



Customers should certainly have the option of “opting out” of having a meter on their house.

But we do not feel that an “opt out” policy is a solution to the public health problem.

You can’t “opt out” of all the meters on the wall of your rental apartment complex. Or the ones on the wall of the complex right across the alley from your apartment.

You can’t “opt out” of exposure to your neighbor’s meter, that is ten feet away from your bedroom window.

You can’t “opt out” of exposure to the meter on the other side of your bedroom wall if you are a baby in a crib.

You can’t “opt out” of exposure to transmissions from the radio tower 100 meters from your house.

An “opt out” doesn’t protect people who DON’T opt out, if a health hazard is being produced.

Strategies for solving the problem:

Residential exposures to constant RF transmissions are harmful, even at low levels, and should be avoided.

A better solution is to recognize the potential problem, and design a system to truly minimize risk, by reducing transmissions to a functional minimum.

Constant transmissions can and should be avoided.

Strategies for solving the problem:

Frequent and repetitive intermittent transmissions are also probably harmful, and should be avoided.

MESH network meters like the ones EWEB was testing a year ago are a case in point, as they transmit multiple times a minute. **EWEB staff should be applauded for steering away from that technology.**

Our group thinks that the MESH technology in the in-home “Zigbee” network is also likely to be harmful to the public. It is lower powered, but contemporary research shows that chronic low power RF exposures can have serious consequences.

It would be a mistake for EWEB to get married to this technology, and invest many millions of dollars in an infrastructure that will become less and less acceptable to the public down the line.

The solution here (for the smaller percentage of your customers who may want to share this functionality in the future) might be to **explore a broadband communication option** that allows them to remotely adjust their thermostat over the internet, rather than via the smart meter network.

80% of the households in our area have broadband at the present time.

Strategies for solving the problem:

Nocturnal exposures are more problematic than daytime exposures.

RF's potential to suppress nocturnal melatonin secretion and disturb sleep is a serious problem.

Chronic suppression of melatonin can increase incidence of breast cancer and some other cancers.

The best solution is to avoid nocturnal RF transmissions.

Fortunately, EWEB doesn't need real-time access to billing data. So **it shouldn't be necessary to upload usage data at night.**

A broadband remote control option would eliminate the need for tower -> meter communications at night.

Strategies for solving the problem:

Occasional and infrequent daytime exposures are much less likely to cause an increase in chronic problems for the population at large.

Industry routines are to upload usage data every four hours. But EWEB is only billing once a month.

The solution here is to upload usage data much less frequently.

If you can fit all the Beatles albums plus Bach's Mass in B Minor on an iPod, it should be possible to design a smart meter protocol that only requires an upload once every two weeks, or less frequently.

This is the direction the industry needs to go in.

EWEB should explore this issue with their vendors, and push for this sort of functionality in any system that they purchase.

Strategies for solving the problem:

Occasional and infrequent daytime exposures may still provoke acute symptoms in a portion of the population.

People with EHS will be likely to react acutely to intermittent exposures in their residential neighborhoods, even if these exposures are infrequent.

Part of addressing this problem is to upload data as infrequently as possible.

A data upload by the system once every two weeks might not be a significant release in exposure burden to this part of the population, where a data upload every four hours could be expected to have significant adverse consequences on their quality of life.

Infrequent transmissions at predictable times might be much easier for these people to deal with.

They will certainly want to opt out of having a meter on their house.

Other steps might also be found to mitigate residential exposures for these members of the population, if RF communications are to be used by EWEB.

Finding a “smart” way to move forward:



We're all trying to work towards the benefit of our community and our environment.

The AMI program is an attempt to reduce EWEB's carbon footprint, aid in energy conservation, and enable more effective utilization of renewable energy sources down the line.

These are laudable goals.

The key thing here is to avoid solving one problem with tools that create a new one.

If EWEB is really “smart” about how we use “smart meters”, the advisory group thinks that it should be possible to find a way to make it work without harming the community.

This will require a progressive approach that looks at some new ways to approach the engineering problem.