Acute Therapies at University District Oregon Rehabilitation Center at University	versity District - Phone:	(541) 686-7363
Outpatient Rehabilitation Services a	t University District - Pl	none: (541) 686-7085
Patient Name: MIKC	STOOKS	5/29
The results of your Modified Barium	Swallow / In-Clinic	Swallow Evaluation onshowed
It is recommended that you follow the solution of the solution	hese diet recommen	dations:
() Nothing orally (NPO)	() Dysphagia Adva	anced
() Dysphagia Puree	() Mechanical Soft	
) Dysphagia Transitional	Regular (genera	1) small bites
-IQUIDS:		
) Nothing orally (NPO)	() Ice chips only	11
Thin (water, coffee, tea, Ensure,	all juices) SM	Il sips
) Nectar-thick (buttermilk, nectars) Honey-thick (can't drink it throug	, iomato, milksnakes	, bienderized soups, cream) on't need a spoon)
) Pudding-thick (pudding, custard		
OMPENSATORY STRATEGIES:		100 million
	Cupandalan	
ositioning:	Supervision:	Medication:
Upright at 90 degrees	Supervision: () Intermittent	Medication: () With water
Upright at 90 degrees) Chin-tuck positioning	() Intermittent() Constant	() With water
Upright at 90 degrees) Chin-tuck positioning) Turn head right/left	() Intermittent() Constant() Feeding needed	 () With water In pureed food () Crushed in pureed
Upright at 90 degrees) Chin-tuck positioning) Turn head right/left) Tilt head right/left	() Intermittent() Constant	 () With water In pureed food () Crushed in pureed
Upright at 90 degrees) Chin-tuck positioning) Turn head right/left) Tilt head right/left) Other:	 () Intermittent () Constant () Feeding needed () Feeds self with s	 () With water In pureed food () Crushed in pureed () With thick liquid
 Upright at 90 degrees Chin-tuck positioning Turn head right/left Tilt head right/left Other: or A Safe Swallow Please Take small bites/sips 	 () Intermittent () Constant () Feeding needed () Feeds self with s () None 	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow
 Upright at 90 degrees Chin-tuck positioning Turn head right/left Tilt head right/left Other: or A Safe Swallow Please Take small bites/sips Wait for each swallow before tak 	 () Intermittent () Constant () Feeding needed () Feeds self with s () None 	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow () Remain upright approximately
 Upright at 90 degrees Chin-tuck positioning Turn head right/left Tilt head right/left Other: or A Safe Swallow Please Take small bites/sips Wait for each swallow before tak Alternate food/liquid 	 () Intermittent () Constant () Feeding needed () Feeds self with s () None 	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow () Remain upright approximately 30-45 minutes after eating.
 Upright at 90 degrees Chin-tuck positioning Turn head right/left Tilt head right/left Other: or A Safe Swallow Please Take small bites/sips Wait for each swallow before tak 	 () Intermittent () Constant () Feeding needed () Feeds self with s () None 	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow () Remain upright approximately
 Upright at 90 degrees Chin-tuck positioning Turn head right/left Tilt head right/left Other:	 () Intermittent () Constant () Feeding needed () Feeds self with s () None 	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow () Remain upright approximately 30-45 minutes after eating. () If coughing occurs, stop eating () Do not feed patient if tired or sleepy Swallow first, then talk
 Upright at 90 degrees Chin-tuck positioning Turn head right/left Tilt head right/left Other: or A Safe Swallow Please Take small bites/sips Wait for each swallow before tak Alternate food/liquid Use a straw Do not use a straw Check for pocketing food Rinse mouth/dentures after eatin 	 () Intermittent () Constant () Feeding needed () Feeds self with s () None 	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow () Remain upright approximately 30-45 minutes after eating. () If coughing occurs, stop eating () Do not feed patient if tired or sleepy
 Upright at 90 degrees Chin-tuck positioning Turn head right/left Tilt head right/left Other: or A Safe Swallow Please Take small bites/sips Wait for each swallow before tak Alternate food/liquid Use a straw Do not use a straw Check for pocketing food Rinse mouth/dentures after eatin Clear throat then swallow if gurgl 	 () Intermittent () Constant () Feeding needed () Feeds self with s () None 	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow () Remain upright approximately 30-45 minutes after eating. () If coughing occurs, stop eating () Do not feed patient if tired or sleepy Swallow first, then talk
 Upright at 90 degrees Chin-tuck positioning Turn head right/left Tilt head right/left Other: or A Safe Swallow Please Take small bites/sips Wait for each swallow before tak Alternate food/liquid Use a straw Do not use a straw Check for pocketing food Rinse mouth/dentures after eatin 	 () Intermittent () Constant () Feeding needed () Feeds self with s () None 	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow () Remain upright approximately 30-45 minutes after eating. () If coughing occurs, stop eating () Do not feed patient if tired or sleepy Swallow first, then talk
 Upright at 90 degrees Chin-tuck positioning Turn head right/left Tilt head right/left Other: or A Safe Swallow Please Take small bites/sips Wait for each swallow before tak Alternate food/liquid Use a straw Do not use a straw Check for pocketing food Rinse mouth/dentures after eatin Clear throat then swallow if gurgl Other: Follow-up with the Speech-Langu 	 () Intermittent () Constant () Feeding needed () Feeds self with s () None ing another bite/sip g y voice Jage Pathologist is response 	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow () Remain upright approximately 30-45 minutes after eating. () If coughing occurs, stop eating () Do not feed patient if tired or sleepy Swallow first, then talk () Place food right / left / center of mouth
 Upright at 90 degrees Chin-tuck positioning Turn head right/left Tilt head right/left Other:	 () Intermittent () Constant () Feeding needed () Feeds self with s () None ing another bite/sip g y voice Jage Pathologist is response 	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow () Remain upright approximately 30-45 minutes after eating. () If coughing occurs, stop eating () Do not feed patient if tired or sleepy Swallow first, then talk () Place food right / left / center of mouth
 Upright at 90 degrees Chin-tuck positioning Turn head right/left Tilt head right/left Other: or A Safe Swallow Please Take small bites/sips Wait for each swallow before tak Alternate food/liquid Use a straw Do not use a straw Check for pocketing food Rinse mouth/dentures after eatin Clear throat then swallow if gurgl Other: Follow-up with the Speech-Language 	 () Intermittent () Constant () Feeding needed () Feeds self with s () None ing another bite/sip g y voice Jage Pathologist is response 	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow () Remain upright approximately 30-45 minutes after eating. () If coughing occurs, stop eating () Do not feed patient if tired or sleepy Swallow first, then talk () Place food right / left / center of mouth
 Upright at 90 degrees Chin-tuck positioning Turn head right/left Tilt head right/left Other: or A Safe Swallow Please Take small bites/sips Wait for each swallow before tak Alternate food/liquid Use a straw Do not use a straw Check for pocketing food Rinse mouth/dentures after eatin Clear throat then swallow if gurgl Other: Follow-up with the Speech-Languappointment will be scheduled after 	 () Intermittent () Constant () Feeding needed () Feeds self with s () None ing another bite/sip g y voice uage Pathologist is reter we receive orders	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow () Remain upright approximately 30-45 minutes after eating. () If coughing occurs, stop eating () Do not feed patient if tired or sleepy Swallow first, then talk () Place food right / left / center of mouth
 Upright at 90 degrees Chin-tuck positioning Turn head right/left Tilt head right/left Other:	 () Intermittent () Constant () Feeding needed () Feeds self with s () None ing another bite/sip g y voice Jage Pathologist is reter we receive orders with MS	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow () Remain upright approximately 30-45 minutes after eating. () If coughing occurs, stop eating () Do not feed patient if tired or sleepy Swallow first, then talk () Place food right / left / center of mouth
) Turn head right/left) Tilt head right/left) Other: or A Safe Swallow Please Take small bites/sips Wait for each swallow before tak) Alternate food/liquid) Use a straw Do not use a straw) Check for pocketing food Rinse mouth/dentures after eatin) Clear throat then swallow if gurgl Other: Follow-up with the Speech-Language appointment will be scheduled after () () () () () () () () () () () () () (() Intermittent () Constant () Feeding needed () Feeds self with s () None ing another bite/sip g y voice Jage Pathologist is reter we receive orders with MS	 () With water In pureed food () Crushed in pureed () With thick liquid 1 pill at a time Swallow twice: dry swallow () Remain upright approximately 30-45 minutes after eating. () If coughing occurs, stop eating () Do not feed patient if tired or sleepy Swallow first, then talk () Place food right / left / center of mouth